



Red Listed medicinal plants of South Africa: Status, trends, and assessment challenges

V.L. Williams^{a,*}, J.E. Victor^b, N.R. Crouch^{c,d}

^a School of Animal Plant and Environmental Sciences, University of the Witwatersrand, Private Bag 3, Wits 2050, South Africa

^b National Herbarium, SANBI, Private Bag X101, Pretoria 0001, South Africa

^c Ethnobotany Unit, SANBI, PO Box 52099, Berea Road, 4007, South Africa

^d School of Chemistry and Physics, University of KwaZulu-Natal, 4041 Durban, South Africa

ARTICLE INFO

Article history:

Received 8 March 2012

Received in revised form 22 September 2012

Accepted 24 January 2013

Available online 28 February 2013

Edited by B-E Van Wyk

Keywords:

Extinction

GSPC

IUCN

Red List Index

South Africa

Threatened species

Traditional medicine

ABSTRACT

In 2009, South Africa completed the IUCN Red List assessments of 20,456 indigenous vascular plant taxa. During that process, medicinal plant species (especially those sold in informal *muthi* markets) were identified so that potential extinction risks posed to these species could be assessed. The present study examines and analyses the recently documented threat statuses of South African ethnomedicinal taxa, including the number of species used, revealing family richness and the degree of endemism, and calculates the Red List Index (RLI) of species survival to measure the relative degree of threat to medicinal species. Approximately 2062 indigenous plant species (10% of the total flora) have been recorded as being used for traditional medicine in South Africa, of which it has been determined that 82 species (0.4% of the total national flora) are threatened with extinction at a national level in the short and medium terms and a further 100 species are of conservation concern (including two species already extinct in the wild). Thirty-two percent of the taxa have been recorded in traditional medicine markets in the provinces of KwaZulu-Natal, Gauteng, Eastern Cape, Mpumalanga and Limpopo. The study also reflects on the challenges associated with Red List evaluations of medicinal species, many of which, based on market reports, are extracted at a seemingly unsustainable rate. In contrast to the majority of species enumerated in the Red List of South African plants, medicinal taxa are often widespread, with large extents of occurrence. Accordingly, the population decline criteria have necessarily been applied to assess threats to their existence, even though accurate figures for numbers of remaining individuals, areas of occupancy, quantities harvested, and regeneration times are often found lacking. Factors leading to susceptibility of plant species to extinction as a result of harvesting pressure are discussed. The current findings reveal a need for greater emphasis on focussed population level research on prioritised medicinal plant species.

© 2013 SAAB. Published by Elsevier B.V. All rights reserved.

1. Introduction

Despite the availability and accessibility of western medicine in the 21st century, a wide range of cultural communities in South Africa still depend on and often prefer traditional medicine (TM) or '*muthi*' as an important component of primary health care. It was estimated that 72% of Black South Africans, even in urban areas, subscribe to traditional health care systems involving the consumption of medicinal plants, and that more than 70,000 tonnes of plant material is consumed in South Africa each year, with at least 134,000 income-earning opportunities generated by the trade in medicinal plants and related products (Mander et al., 2007). Medicinal plants are traded through several large markets throughout South Africa, particularly in the summer-rainfall region (Mander, 1997; Mander, 1998; Botha et al., 2001; Dold and

Cocks, 2002; Von Ahlefeldt et al., 2003; Williams, 2003, 2007). Much of the material supplied is obtained mainly from harvesting wild plant resources, leading to concerns over sustainability. Threats to the resource base are further compounded by habitat transformation occurring over parts of the country (Fairbanks et al., 2000), with some vegetation types more adversely affected than others (Mucina and Rutherford, 2006). The rapidly urbanising landscape of Gauteng, for instance, requires medicinal plant users based on the Witwatersrand to source their plants from increasingly further afield (Williams, 2003).

With the population of South Africa continuing to increase (Statistics South Africa, 2007) in the midst of an HIV/AIDS epidemic that requires the extensive herbal treatment of opportunistic infections, the demand for medicinal plant resources is growing apace. A wide range of species are already showing signs of unsustainable harvesting, with the size of the traded components (e.g. bulbs) decreasing, distances to harvesting source increasing (Mander, 1998), supply becoming increasingly irregular and some plants becoming unavailable in certain markets (Te Beest, 2003). There is also documented extirpation at local (e.g. *Warburgia salutaris*) (Gerstner, 1938; Mander, 1998; Botha et al., 2004a) and provincial (e.g. *Siphonochilus aethiopicus*) (Crouch et

* Corresponding author. Tel.: +27 117176336.

E-mail address: vivwill@netdial.co.za (V.L. Williams).

al., 2000) levels. Only one species (*Encephalartos woodii*) is known to be extinct in the wild as a result of over-harvesting for medicine, having been on the brink of extinction at the time of its scientific discovery at the turn of the 19th Century (Crouch et al., 2003a). Unsustainable medicinal plant use in South Africa is, however, not a recent concern. As early as 1946, Father Jacob Gerstner, a Zululand missionary, anticipated the extinctions of “doomed” plant species used for traditional medicine (Gerstner, 1946). He accordingly recommended their cultivation by state nurseries, a relevant suggestion acted on by neither national nor provincial governments of the time. Gerstner (1946) reported witnessing 40 to 50 bags of *Erythrophleum lasianthum* (Fabaceae) being railed to Durban on a single day, presumably to supply *muthi* markets.

There are three large, permanent urban *muthi* markets in South Africa requiring a steady supply of plant material: the Warwick Triangle and Ezimbuzini markets in Durban, and the Faraday market in Johannesburg. Permanent street traders and ephemeral pension day markets, selling medicinal plants, are also present in smaller urban centres throughout the country (Dold and Cocks, 2002; Botha et al., 2004b). In addition to informal street markets are formal *muthi* stores found in almost all urban areas in the region. Less attention has been focussed on *muthi* stores with regard to their contribution to health care services and medicinal plant exploitation, but work in the Eastern Cape and the Witwatersrand shows that a large number of medicinal plant species are traded in these stores (Cocks, 1996; Williams et al., 2001).

The Species Survival Commission of the IUCN devised an objective system of assessing extinction risks based on a set of five quantitative criteria (A–E) with threshold values that delimit the various categories of threat. The IUCN Red List Categories and Criteria Version 3.1 (2001) were used in the assessments of 20,456 indigenous South African vascular plant taxa (Raimondo et al., 2009a; Von Staden et al., 2009a), the first time in the world that a country's entire flora has been assessed. The potential extinction risks posed to plants, including those threatened by harvesting for the medicinal plant trade, were assessed by means of the quantitative criteria, along with other considerations (e.g. habitat loss), to determine their Red List status. Assessing the conservation status of widespread plants, in which category many heavily traded medicinal taxa fit does, however, pose unique challenges. Not only is it difficult to accurately estimate a percentage decline over a set period of time (required if taxa are listed under the population decline Criteria A and/or C), but often species undergoing a continuing decline are abundant and widespread. Accordingly, they do not qualify for the restricted-range Criteria B and/or D. Furthermore, while many medicinal species may be facing a much lower risk of extinction, they still require conservation action to diminish future risks of regional or localised extirpations. In order to meet local conservation concerns for species that are declining but not necessarily in danger of extinction, Victor and Keith (2004) developed additional categories (Critically Rare, Rare and Declining) to flag taxa of conservation concern that would be classified as Least Concern (LC) according to the IUCN system. While these categories are only applied in South Africa, they are pertinent for ethnomedicinal taxa that should be considered in conservation prioritisation processes to ensure that they do not become threatened in the future should deterministic factors (e.g. over-exploitation) in their extinction risks persist.

A challenge for assessing the Red List statuses of medicinal taxa is the availability and accessibility of sufficient quantitative and qualitative data to determine the extent of population decline; however, inferences can be made from observing trends regarding individual plant sizes in *muthi* markets. For instance, Williams et al. (2007a,b) considered market-derived data as a surrogate in an attempt to assess the condition of wild standing stocks. They found statistically significant decreases in the market availability of large bulbs or thick tree bark over time to signal both population decline within a species' area of occupancy, and the reduced occurrence of larger, reproductive individuals. In 1994, for example, *Bowiea volubilis* (Hyacinthaceae) bulbs with a diameter > 15 cm were frequently encountered in Johannesburg *muthi* shops. By 2007,

however, 93% of *B. volubilis* bulbs recorded in the Faraday market were <4 cm diameter (Brueton, in preparation). These market data, where available, in conjunction with expert observations, were used in the course of assessing the Red List status of medicinal species known to be traded.

In light of South Africa's significant achievement in assessing the threat statuses of its entire flora, and the unique availability of a complete set of baseline information by which one can fully compare groups of flora, in this paper we analyse and reflect on the recently documented threat and conservation statuses of South African medicinal plants. The objectives were to examine patterns and trends in the taxa used, and to discuss factors leading to the susceptibility of plant species to decline as a result of harvesting pressures. Since medicinal plants are an important resource to the majority of South Africans, the information presented should inform decisions taken by policy makers within the domain of biodiversity conservation and sustainable livelihoods.

2. Methods

2.1. Species used and Red List assessments

In order to determine the number of threatened medicinal plants and their conservation status, a list of plant species cited as being used and/or traded for traditional medicine was compiled. The primary source of this information was Arnold et al. (2002) and the references cited therein, including tertiary literature reports such as Watt and Breyer-Brandwijk (1962) and Hutchings et al. (1996). However, Arnold et al. (2002) list plants that occur in southern Africa and that have been cited as being used for traditional medicine anywhere else, though most of them have been employed in the subregion. Since this paper is about South African ethnomedicinals only, the plants were checked against the references and only species that occur in South Africa and that were cited as being used/traded in South Africa were retained from the original Arnold et al. (2002) checklist. A further comprehensive literature search was conducted through published material to justify our decision on what species to retain. More than 140 species were further added to our checklist of medicinal plants from the following references: Anonymous (c.1994), Brandt et al. (1995), Crouch and Hutchings (1999), Crouch and Krynanuw (1999), Crouch et al. (2005a,b), Cousins et al. (2012), Dold and Cocks (2000, 2001, 2002), Dzerefos and Witkowski (2001), Hanekom (1967), Kamatou et al. (2008), Kroon (1999), Lourens et al. (2008), Loxton, Venn and Associates (1994), Makgakga (1995), Makunga et al. (2008), Matsiliza (1997), McGraw et al. (2008), More et al. (2008), Netshiungani (1981), Newton (1993), Pooley (1993, 1998), Scott-Shaw (1999), Smith and Crouch (1995), Sobiecki (2002), Van Vuuren (2008), Van Wyk (2008), Van Wyk et al. (1997), Veale et al. (1992), Verschaeve and Van Staden (2008), Von Ahlefeldt et al. (2003), Williams (2003, 2007). The species list is, however, not static and papers published subsequent to this research will add to our knowledge on the number of species used and traded for traditional medicine in South Africa. For example, Moeng and Potgieter (2011) listed two species not captured in our list as being sold in shops and markets in the Limpopo Province. For the purpose of this paper, however, new species recorded in literature published after February 2010 were not taken into consideration.

Not all plant species that are used for traditional medicine are commercially traded and threatened by destructive and unsustainable harvesting practices. Hence, when the conservation statuses of ethnomedicinals were assessed according to the IUCN Red List categories and criteria, a distinction was made between plant species recorded in trade in the medicinal markets of KwaZulu-Natal, Gauteng, Eastern Cape, Mpumalanga and Limpopo Provinces, and those plant species that were cited as being used only but have yet to be identified in the markets. Special attention was also paid to the most heavily traded species, since the likelihood of these plants being threatened by the medicinal plant trade is high (Williams, 2007). Given that many heavily traded medicinal taxa

are widespread and do not qualify under the restricted-range Criteria B and D, an extensive literature search and a workshop with eight ethnobotanists was conducted to try and obtain information to support a listing under the population decline Criteria A and C (Von Staden et al., 2009b). The detailed information obtained during this process sets the medicinal plant assessments apart from the conservation assessments of other South African plants (Von Staden et al., 2009b).

The information collected to support a threat assessment based on population decline as a result of harvesting for the informal plant trade includes: observed losses of individuals or subpopulations over time; the extent of the decline in the distribution range, mainly inferred through information on past and present distributions; relative abundance (e.g. locally frequent, or uncommon in the field); the quantities of plant material traded (Cunningham, 1988; Mander, 1998; Williams, 2007; Williams et al., 2007a); the type of harvesting (i.e. destructive vs. non-destructive); life cycle characteristics (e.g. reproductive capabilities and growth rate); and popularity in the *muthi* markets (Cunningham, 1988; Mander, 1998; Williams, 2007). In addition, information on suspected levels of population decline due to trade was estimated and extrapolated from observed losses of subpopulations, as observed by fieldworkers. These data were also sourced from ethnobotanists during the course of the expert Red Listing medicinal plant workshop (Von Staden et al., 2009b); participants reported on observed changes with time in the abundance, availability and size of individual plants and plant parts. While the expert workshop for ethnomedicinals looked mainly at popular plants (which were invariably those in trade), post-workshop research elucidated further non-trade related threats to some species e.g. habitat loss to commercial agriculture for *Stangeria eriopus*. These were taken into account in the assessments of the relevant species. Since harvesting is usually not a threat to non-traded medicinal taxa, evaluations of the threats to these species were mostly reliant on information contributed by many other knowledgeable botanists who assisted with the Red Listing process.

The assessment of the conservation status of medicinal plant species was carried out using the IUCN Red List Criteria Version 3.1 (2001). There are five broad criteria (A to E) based on quantitative thresholds that are used to assess the probabilities of extinction. If a species does not meet any of the criteria, it is classified as Least Concern (LC). Taxa can be classified into one of the three categories of threat, i.e. Critically Endangered (CR), Endangered (EN) or Vulnerable (VU); or else they are placed into Near Threatened (NT), Data Deficient (DD), Extinct (EX) or Extinct in the wild (EW). For Criterion A, the taxon must show a population reduction (measured as a percentage) over a set time period. Taxa qualifying under Criterion B would have a limited geographic range or restricted area of occupancy and at the same time meet two of the following three subcriteria: a continuing decline; a severely fragmented population or fewer than a threshold number of locations; extreme fluctuations in specified population parameters. Taxa with small global populations would qualify for a category of threat under Criterion C if the total population size is less than a threshold number of mature individuals and there is either a continuing decline over a set period or a continuing decline in numbers of mature individuals combined with small or single subpopulations and/or extreme fluctuations in the number of mature individuals. Taxa with very small and/or very restricted populations, and that are potentially subject to threatening processes, would qualify under Criterion D. Where a quantitative analysis indicates a probability of extinction in the wild to be at least 10% within 100 years, a taxon would qualify for a category of threat under Criterion E. For further more detailed explanation and definitions, refer to IUCN (2001). A species classified as LC can additionally be flagged as being of conservation concern either as Rare, Critically Rare or Declining (Victor and Keith, 2004; Von Staden et al., 2009c).

The Red List assessments for the medicinal taxa analysed in this paper are correct as of February 2010. Subsequent changes to the status of seven medicinal species (one status upgrade and six downgrades)

were found when the updated Red List of South African plants version 2012.1 (SANBI, 2012) was announced in May 2012. However, we have not accordingly adjusted the figures and analyses since most changes affect species that are not threatened. Status changes are indicated in the footnotes of Appendix A.

2.2. Red List Index

To evaluate the overall extinction risks to medicinal plant species as a group, we calculated the IUCN Red List Index (RLI) of species survival (Butchart, 2008). The RLI uses information from the Red List to monitor trends in the proportion of species expected to remain extant or to become extinct in the near future (Butchart, 2008), and can be used to explore trends resulting from the impact of specific threats (e.g. unsustainable use) to groups of species (Bubb et al., 2009). While the index is usually calculated as a trend that requires all species in the group to have been assessed for the Red List at least twice (Bubb et al., 2009), RLI values can also be calculated at a single point and the values indicate the relative degree of threat to species in a group at that point in time (S.H.M. Butchart, pers. comm., 2011).

Guidance on the national and regional uses of RLIs is described in Bubb et al. (2009) and Butchart (2008). For the purposes of this study, the RLI was calculated by: A) multiplying the number of species in each Red List category by a category weight (Extinct in the Wild = 5; Critically Endangered = 4; Endangered = 3; Vulnerable = 2; Near Threatened = 1; Least Concern = 0) (Data Deficient and NE taxa are excluded from the analyses since the calculation requires species to have been assessed); B) summing the product of the category scores, and dividing this by the sum of the maximum possible product (i.e. the number of species multiplied by the maximum weight, i.e. 5); and then C) subtracting this value from 1 (Bubb et al., 2009). Since the category of Declining was developed specifically for the South African plant conservation context (Victor and Keith, 2004; Von Staden et al., 2009c) and is not a status recognised at the global level, all Declining species, for the purposes of the RLI, were treated as Least Concern. RLI values closer to 1 indicate that most species are categorised as Least Concern and are expected to remain extant in the near future without additional conservation action, whereas RLI values closer to zero indicate that a high proportion of species have become extinct or are highly threatened (Bubb et al., 2009; Butchart, 2008). Since 80% of South Africa's flora was assessed for the first time in 2009 (Raimondo et al., 2009a), it was not possible to derive RLI values for most species from the previous South African Red List Assessment (Victor, 2002); hence, temporal trends in the extinction risks to South African medicinal plants could not be measured. The RLI was calculated for the entire list of medicinal plants ($n=2000$ species; EW, CR, EN, VU, NT and LC species only), and then the data were divided into two subgroups and compared accordingly, namely species that are traded ($n=647$ species) versus species that are not known to be traded ($n=1353$ species).

It is important to note that, with the exception of endemic species, the Red List statuses used in the calculation of the RLIs for the medicinal plants are all, by default, national assessments because the South African Red List assessments apply only to the section of the global population that occurs within South Africa's borders (Von Staden et al., 2009c). For endemic species, however, the national status is the same as the global status. Hence, when interpreting the RLI values in this paper at a national level, it cannot be assumed that the levels of threat to a group of species are indicative of the survival expectancies at a global level, since the proportion of more widespread non-endemic species must also be considered.

In addition to the RLI, we examined threats at plant family level and calculated the percentage of medicinal species within broad threat categories that were endemic or not endemic to South Africa and the FSA (Flora of Southern Africa) region.

3. Results and discussion

3.1. Patterns and trends in the Red List of South African medicinal plants

There are over 3400 plant species in southern Africa that are used for medicinal purposes (Arnold et al., 2002); of these, 2062 taxa have specifically been cited as used and/or traded in South Africa (10.1% of the total national flora). Furthermore, 32% of South Africa's plant ethnomedicines have been recorded in *muthi* markets in KwaZulu-Natal, Gauteng, Eastern Cape, Mpumalanga and Limpopo (Table 1). Overall, 9.2% were assessed according to threats driven by harvesting for the traditional medicine trade (Table 1). Although the Red List is not static and the threat statuses of some species will change in future to reflect both population and plant use dynamics, there are currently two medicinal plant species Extinct in the Wild and 82 species (4% of the medicinal flora, 0.4% of the national flora) threatened with extinction at a national level (Table 1; Fig. 1). Fourteen species are Critically Endangered, 19 Endangered and 49 Vulnerable (Table 1). A further 100 species are of conservation concern, of which 37 (1.8%) are Near Threatened, 36 Declining, four are Data Deficient (DD) as they are suspected to be threatened but insufficient information is available to place them in a category of threat, and 21 species are either Rare or Critically Rare. The 37 Near Threatened and 36 Declining species (the latter representing 69% of the total number of 52 Declining species assessed in South Africa) are not currently facing a high risk of extinction, but may very well do so in the near future if threats persist. Taxa that are threatened and of conservation concern account for 8.8% of the assessed species (Fig. 1); hence the majority of ethnomedicinals were assessed as Least Concern (88.4%). Appendix A lists the species that are threatened and/or of conservation concern.

Fifty percent or more of the taxa within each threatened category or category of conservation concern have been recorded in the *muthi* markets, especially the Declining (94%), Near Threatened (81%) and Vulnerable (74%) taxa (Table 1); furthermore, 68% of the 82 threatened medicinal species are traded. Hence, traded species are more threatened than non-traded species and persistent commercial exploitation could result in the future upgrading of the Red List statuses of several species that are currently at risk. Less threatened by the traditional medicine trade, however, are Least Concern taxa and only 29% of the 1822 taxa have been recorded in the markets. Threats posed by the traditional medicine trade were specifically used in the assessments of 9.2% of the total number of ethnomedicinal species (Table 1, column 6); within each Red List category, the percentage of species recorded in trade and the percentage of species assessed

Table 1
Classification of medicinal plant species in South Africa according to IUCN Red List categories of threat^a.

National Red List status category ^b	Total no. species	No. recorded in trade	% recorded in trade	No. assessed as a traded medicinal	% assessed as a traded medicinal
EW	2				
CR	14	7	50.0%	7	50.0%
EN	19	13	68.4%	12	63.2%
VU	49	36	73.5%	35	71.4%
NT	37	30	81.1%	26	70.3%
Declining	36	34	94.4%	33	91.7%
Rare	21	5	23.8%	0	0
DDD	4	3	75.0%	1	25.0%
LC	1822	521	28.6%	70	3.8%
DDT	17	6	35.3%	5	29.4%
NE	41	1	2.4%		
Total	2062	656	31.8%	189	9.2%

^a Data are correct as of February 2010; for the most recent statuses see <http://redlist.sanbi.org>.

^b Abbreviations: EW = Extinct in the Wild; CR = Critically Endangered; EN = Endangered; VU = Vulnerable; NT = Near Threatened; Declining = LC species declining; Rare = LC species considered Critically Rare or Rare; LC = Least Concern; DDD = Data Deficient–Insufficient Information; DDT = Data Deficient–Taxonomically Uncertain; NE = Not Evaluated.

based on their class as a traded medicinal are similar, except for LC species where the majority of species are not traded (Table 1, cf. columns 4 and 6). For example, of the 49 Vulnerable taxa (Table 1), 36 were recorded in the markets and the Red List assessments of 35 of these were based on population decline information derived and inferred from ethnobotanical literature and market data. Taxa not evaluated according to threats posed by harvesting for the TM trade were thus evaluated according to other factors that pose a more significant threat to their persistence in the wild.

As has been previously stated, a challenge with assessing medicinal plants according to IUCN criteria is that they are typically widespread, occur in hundreds of locations, and therefore do not qualify for listing under the restricted-range Criteria B and D. To verify this observation, we analysed the proportion of threatened and Near Threatened taxa that were listed according to Criteria B and D versus Criteria A and C for 119 medicinal and 2762 South African plants assessed in 2009 (derived from Raimondo et al., 2009a). Currently on the IUCN Red List, 80% of plants assessed in the last 10 years are listed based on the range estimate criteria B and D2 (Rivers et al., 2011). In South Africa, the percentage is similar, viz. 81% (Table 2). For medicinal species, however, we found that of the 119 threatened and Near Threatened taxa, 44% were listed according to Criteria A and/or C, 41% were listed according to Criteria B and/or D, and 14% were listed according to B with A and/or C (Table 2) – hence, 55% of all medicinal taxa are listed based on range as risk factor. However, when one looks more closely at the species recorded in the *muthi* markets that were assessed according to the risks posed by the trade, we find that 53% were listed according to the population decline criteria and only 31% could be listed according to the restricted-range criteria (Table 2). Threatened and Near Threatened medicinal species that were *not* recorded in the *muthi* markets were usually listed according to Criteria B and/or D (67%) compared to Criteria A and/or C (18%) (Table 2). Hence, use and harvesting pose less of a conservation threat to non-traded medicinals relative to their current limited distribution range. The impact on specific taxa by local communities in antiquity is for the most part indiscernible, although early ethnographers have alluded occasionally to plant extinction events as a result of unsustainable utilisation (Dicke, 1926).

Obtaining information to support population decline is not as easy as calculating the Extent of Occurrence (EOO, the quantitative criterion B1). Population decline data is seldom found in publications, hence the knowledge of experts was vital in accomplishing the medicinal plant assessments. Had this knowledge not been obtained, then it would have been near-impossible to assign a threat status to the majority of popularly traded medicinal plants, since their widespread distributions would have placed them outside of the quantitative thresholds for listing as threatened or Near Threatened according to B1.

3.2. Endemism

Overall, levels of endemism are low across South Africa's 2062 medicinal species (27% endemic, Fig. 2), reflecting generally widespread distributions, whereas the majority of threatened (CR, EN, VU) and Near Threatened medicinal species are endemic (68% and 57% respectively) (Fig. 2). By contrast, 72% of LC species and only 18% of threatened species are not endemic. Accordingly, medicinal plant species endemic to South Africa and the FSA region are more likely to be threatened.

3.3. Plant family trends

The 2062 taxa used for traditional medicine are from 171 families, the largest of which is the Asteraceae (12.5% of the total number of South African species; for context of scale, the size of the family in this country is provided) (Table 3; Appendix B). The next largest family is the Fabaceae (7.7%) followed by Apocynaceae (5.0%) and Asphodelaceae (4.0%) (Table 3). The eight families with 50 or more medicinal plant species contain 41% of the total number of species used. The first three




Fig. 1. Taxa classified according to the South African Red List Categories as a proportion of the total number of plants used for traditional medicine in South Africa. 'Threatened' includes the IUCN categories Critically Endangered, Endangered and Vulnerable.

families listed in Table 3 were shown by Douwes et al. (2008) to be better represented in the ethnomedicinal flora than in the overall flora of the subregion. Based on regression analyses it has been possible to identify families that are exceptionally well utilised relative to their overall prevalence in the flora (Douwes et al., 2008), and further in the marketplace (Williams et al., 2000). Future Red List assessments of the medicinal flora of South Africa can accordingly be directed, in part, on the basis of such findings.

Families that have the highest proportion of threatened and Near Threatened taxa include Zamiaceae (92% of the 24 species used are threatened or Near Threatened), Amaryllidaceae (46% of 42 species), Hyacinthaceae (20% of 44 species), Dioscoreaceae (38% of 8 species) and Lauraceae (43% of 7 species) (Appendix B). In addition, there are five families represented by one species in the medicinal flora and that species is threatened, namely Canellaceae, Cornaceae, Hydrostachyaceae, Stangeriaceae and Zingiberaceae. These five families are also monotypic at the level of family and genus in South Africa (i.e. represented by a single genus and species respectively) and are hence of notable conservation value.

3.4. Red List Index

The Red List Index value for all the medicinal species was found to be 0.974 ($n = 2000$ species, excluding Data Deficient and NE taxa). This indicates that a high proportion of medicinal taxa are categorised as LC and that 97.4% of the medicinal plants in South Africa are

Table 2

Criteria used to assess the status of threatened (CR, EN, VU) and Near Threatened (NT) medicinal (traded and not-traded) and non-medicinal South African species.

	Population decline criteria A and/or C	Restricted-range criteria B and/or D	Combination of B and/or D with A and/or C
Traded plants ($n = 86$)	53% (40% A only)	31% (22% B only)	15%
Non-traded plants ($n = 33$)	18% (15% A only)	67% (61% B only)	12%
Total medicinals ($n = 119$)	44% (33% A only)	41% (33% B only)	14%
SA flora ($n = 2762$) ^a	9.0%	81%	10%

^a Total number of threatened and Near Threatened plants in South Africa (including 119 medicinal flora) for which criteria A–D were assigned based on the 2009 assessments (Raimondo et al., 2009a). This is not the total number of plants of conservation concern.

expected to remain extant in the near future without additional conservation intervention, and without considering the impact of climate change, further habitat loss, or negative trends in harvesting patterns. Therefore, in terms of the RLI, only 2.6% of all medicinal taxa are expected to be highly threatened at a national level (this figure differs from the 4% that are actually threatened based on Table 1). However, since only 27% of all medicinals are endemic to South Africa (Fig. 2), the actual proportion of medicinal taxa at risk at a global level will be less than 2.6% (or, <0.3% of the 20,456 indigenous South African flora). Accordingly, although commercial harvesting is a serious concern and is causing significant plant population declines for mainly popular species, the threats posed only affect a very small percentage of plants. However, commercial harvesting is threatening relatively more traded versus non-traded taxa.

To further explore how the exploitation of plants for the traditional medicine trade might be intensifying extinction risks compared to species that are used and have not been recorded in the markets (excluding Data Deficient taxa), two subsets of traded vs. not-traded plants were compared. The RLI values were 0.946 and 0.988 for traded ($n = 647$ species) and not-traded ($n = 1353$) species respectively, indicating that proportionally more plants recorded in the markets are likely to be threatened with extinction in the near future compared to species that are not as exploited. Unsustainable use of medicinal plant resources is not, however, the only threat in South Africa driving population decline for many of these species. Hence, the differences in the RLI values cannot be solely attributed to the impact of harvesting for the trade; the results show how traded species are more threatened than not-traded species by threats that include unsustainable use. When the national Red List of plants is updated in the future, and if the statuses of plants change, then the RLI can be used to show trends in the proportion of medicinal plants used and/or traded for traditional medicine that are being adversely affected by harvesting.

3.5. Geographic and autecological factors

Geographic and autecological factors, in combination with harvesting and utilisation pressures, play a notable role in the susceptibility of plants to population decline and extinction. Species with inherent characteristics such as slow growth and reproductive rates, long generation times and the production of relatively few propagules are rendered especially vulnerable to harvesting and will be more adversely affected than rapidly




Fig. 2. The percentage of medicinal species per threat category that are endemic to SA, endemic to the FSA region, or not endemic.

regenerating species by whatever destructive harvesting practices are exercised by commercial gatherers. Similarly, endemic species with restricted and fragmented distributions are more likely to be extirpated than widespread taxa. However, for medicinal and other utilised plants, these factors need to be considered in conjunction with the desirability/popularity of the species and the amount of harvesting that takes place, as well as the plant part harvested. For example, trees such as *Albizia adianthifolia* (Fabaceae) (LC) tend to tolerate debarking better than more sensitive species such as *Elaeodendron transvaalense* (Celastraceae) (NT A4d), which exhibits poor bark regrowth capacity (Williams and Geldenhuys, 2004).

As taxa with characteristically restricted range distributions, both the Stangeriaceae and Zamiaceae are inherently susceptible to any pressure in that their extremely slow reproductive rate encumbers recolonisation after harvesting, so even small rates of harvesting can be unsustainable. Even seedlings are extracted when encountered, and traded to be grown around homesteads as *intelezi* (protective charm) plants. Although the predominant pressure on cycad populations (particularly for the rarer species) is related to harvesting for horticultural purposes (Raimondo and Donaldson, 2003), the trade in medicinal plant markets also has had a significant impact on several of the more accessible species (Cousins et al., 2011, 2012).

While restricted geographic distributions are a considered factor in the continued existence of about 55% of threatened and Near

Threatened medicinal plants (Table 2), widespread medicinal species are not significantly less at risk of extinction or localised extirpation. We consider two tree species as examples illustrating the complexities of the Red List evaluation process. The endemic Black Stinkwood, *Ocotea bullata* (Lauraceae), for example, has a widespread but somewhat disjunct distribution and an extensive EOO of 659,100 km²; it was evaluated as EN because of reliable quantitative evidence that the population has declined by 53% in the last three generations or 240 years (Williams et al., 2009), largely due to timber and bark exploitation. In IUCN Red List terms, 'generation time' refers to the average age of mature individuals in a population, and *O. bullata* is an example of a highly desirable medicinal species with a long generation length estimated to be a minimum of 80 years (Medicinal Plant Red List Workshop 14/01/2008, SANBI, Durban). Evidence for this generation time was inferred from a) Sim (1906), who estimated that the mean exploitable age of a tree was 80 years old and that the mean annual increment in stem diameter was 6.35 mm; b) data indicating that mean DBH growth rates vary depending on tree size and position with respect to sunlight (Geldenhuys, 2004), and c) that *Ocotea* has long-lived clonal coppice shoots and that some individual trees could be thousands of years old. However, the older a tree is, the less seed production there is and the greater its susceptibility to the root rot pathogen *Phytophthora cinnamomi* (Lubbe and Mostert, 1991). Heavy exploitation of *O. bullata* timber commenced c. 1772 in the Western Cape and extended to Mpumalanga c. 1886 (Fourcade, 1889; Sim, 1906; Phillips, 1931; King, 1939, 1941; Rycroft, 1944; Palmer and Pitman, 1972; McCracken, 1986; Morty and Johnson, 1987). For example, King (1941) estimated that from 1772 to 1938 more than 120,000 m³ of *O. bullata* was felled in forests between George and Plettenberg Bay – an average of 721 m³ per annum. If the average stem yielded 0.4 m³ (Sim, 1906), then an estimate of the total number of trees felled over 167 years is 300,868 – an average of 1802 trees per annum. Many State forests were reduced to a state of complete exhaustion (King, 1939), and in some years commercial exploitation increased to 1255 m³ per annum (total of 3138 trees from 1889 to 1905) (Sim, 1906). Additionally, there has been detrimental exploitation for bark for the traditional medicine trade, documented from c. 1944 (Rycroft, 1944; Taylor, 1961, 1963; Cooper, 1979; Oatley, 1984; Dally, 1984; Morty and Johnson, 1987; Cunningham, 1988; Mander, 1998; Williams et al., 2000; Geldenhuys, 2004). When supporting information for the destructive timber extraction and bark exploitation was

Table 3
Families with 50 or more of the medicinal plant taxa, ranked according to proportion of medicinals.

Family	Number of medicinal species	Percentage of the 2062 taxa in South Africa that are medicinals	Size of the family in the SA flora
Asteraceae	257	12.5%	2236
Fabaceae	159	7.7%	1638
Apocynaceae	104	5.0%	689
Asphodelaceae	83	4.0%	558
Lamiaceae	68	3.3%	264
Euphorbiaceae	65	3.2%	387
Orchidaceae	63	3.1%	487
Malvaceae	50	2.4	332

combined, it was evident that *Ocotea* subpopulations in at least 53% of the quarter-degree squares it once occupied experienced massive declines in tree numbers and density, thereby rendering some of the subpopulations extinct, near-extinct, rare, scarce or fragmented.

By contrast, a tree less widespread in South Africa (*Brackenridgea zanguebarica*, Ochnaceae) has a national EOO of <35 km², and the one known subpopulation in South Africa is significantly disjunct from other African subpopulations occurring northwards to Tanzania (Williams and Raimondo, 2009). The species' restricted distribution is exacerbated by persistent harvesting for its roots and the degradative transformation of a large percentage of its remaining habitat, consequently resulting in an 86% population decline between 1990 and 1997 (Todd, 1999; Todd et al., 2004). This rate of decline is far more rapid than that recorded for *O. bullata*, and *B. zanguebarica* accordingly has a national assessment of CR A2ad; B1ab(ii,v).

The geographic proximity of many medicinal plant species (or their potential substitutes) to each other and in relation to traditional users also plays a large role in their susceptibility to harvesting. Arnold et al. (2002) demonstrated that the documented diversity of ethnomedicinal taxa (per quarter degree square) is most concentrated in northern and eastern South Africa, coinciding with the most fertile and densely settled regions of the country (Statistics South Africa, 2004; Keith and Warren, 2007). *B. volubilis* (Hyacinthaceae) provides a good example of just such a species. This bulbous taxon occurs from the Eastern Cape to Limpopo Province of South Africa, northwards through central Africa to Kenya (Bircher et al., 1998), and grows in a wide variety of habitats ranging from exposed scree to well-shaded forest floors, in the summer-rainfall region (Van Jaarsveld, 1992). *B. volubilis* is under severe pressure from medicinal plant harvesting in the majority of its range in South Africa, with an estimated minimum decline of 30% nationally during the past 30 years (Raimondo et al., 2009b). This estimated decline is based on observations of declines of known subpopulations as well as from measured reductions in the sizes of individual bulbs available in the medicinal plant markets. With a broad extent of occurrence of 486,300 km² (V.L. Williams, unpublished data), exact data on which historically known subpopulations have disappeared are not available. The species is very popular in trade and was rated by herb traders and rural herbalists in KwaZulu-Natal to be among the top six species that have become difficult to source (Cunningham, 1988). In Mpumalanga, only two subpopulations are known: one has been extirpated, and the other (in a private nature reserve) has experienced a 40% decline over the last decade. Other provinces have observed similar trends, with KwaZulu-Natal estimating a decline of 90% (30% during the past 30 years), and in Gauteng, a minimum 60% population decline in the Suikerbosrand Nature Reserve during the past ten years (Raimondo et al., 2009b). The large-scale and unsustainable extraction from nature reserves as well as privately owned land has resulted in a national Red List status of VU A2ad. The current Red List status, however, remains controversial, as some field workers have noted that it reproduces rapidly and is common in the more remote areas of the country (G. Nichols, pers. comm. cited in Raimondo et al., 2009b). Although in itself not quantitatively supported, this criticism may be justified, especially given that no actual figures supporting the percentage decline predicted are available. Inferences have had to be made from overall field observations, and bulb size reduction in the marketplace. In the absence of sufficient conclusive evidence, the precautionary principle is adhered to, and the species therefore remains in the VU category.

Williams (2010) indicated that there is a trend towards threatened medicinal species being more prevalent in a band along the eastern coastline of South Africa, and less threatened medicinal species occurring from the coast to the interior of the country. For these reasons, species outside of the geographic ranges most commonly exploited by harvesters are considerably less likely to be unsustainably targeted by users. The Northern Cape Bokkeveld endemic *Clivia mirabilis* (Hyacinthaceae) (VU D2), for example, is unlikely to be used as much as other members of the genus, all of which are known to be utilised (Crouch et al., 2003b; Williams, 2005)

and are threatened by indiscriminate harvesting practices along the eastern coastline.

Not all species that are popular as medicinal plants and commonly found in the *muthi* markets are threatened with extinction. An example of this is *Elephantorrhiza elephantina* (Fabaceae), a widespread southern African suffrutex occurring in large colonies in grassland and possessing an extensive underground woody rootstock (Brenan, 1970; Schmidt et al., 2002). The species was assessed as Least Concern since there was no evidence to suggest that it is overly threatened by rhizome harvesting despite its popularity in the *muthi* trade (Williams, 2009). *E. elephantina* is heavily harvested, traded and very popular in the Eastern Cape markets (A.P. Dold, pers. comm.) and is fairly common in the Johannesburg and Mpumalanga markets as well (Botha et al., 2001; Williams, 2007). It is believed that the extensive underground stem coppices when damaged and is near-impossible to remove completely (A.P. Dold, pers. comm.), hence lowering the risk of over-exploitation.

4. Conclusions

Medicinal plants are culturally and economically important resources for a large proportion of South Africa's population. Hence, an understanding of their conservation status is important for guiding conservation policy development and action, and contextualizing community-based natural resource management and rural livelihood strategies. Accordingly, we have presented mainly quantitative data that elucidates the state of South Africa's ethnomedicinal taxa in terms of the numbers of species used, the proportion that are traded, and endemic, and threatened. Identified are some of the factors eliciting population declines, and challenges in assessing medicinal species that are mostly widespread compared to similarly threatened non-medicinal plants that tend to be range-restricted. It is hoped that this paper will inform the processes of advising on policy appropriate for the conservation of medicinal plants at national and provincial levels. The data presented are a baseline set against which future quantitative analyses of the conservation status of South Africa's medicinal plants can be compared, on the premise that their status is dynamic. Such comparisons will inform on progress towards achieving specific targets of the 2020 Global Strategy for Plant Conservation (GSPC), most especially Target 2 (an assessment of the conservation status of all known plant species, as far as possible, to guide conservation action) and Target 12 (all wild harvested plant-based products sourced sustainably) (GPPC, 2010).

Factors that influence whether or not taxa will face extinction threats include the size of the distribution range, the relative abundance of mature individuals in the wild and the intensity of the deterministic threat(s). For most medicinally harvested taxa, the extent of their geographic range is less of a factor in predicting risk since most of the species are widespread and the population numbers are still relatively high. However, type of harvesting, popularity and the resultant quantity harvested, as well as inherent life cycle characteristics (autecology) determine the extent of population decline and the sustainability of species use. It is, therefore, anticipated that future Red List updates of ethnomedicinal plants will require further input from experts (as new knowledge becomes available) and the consideration of 'grey' literature sources that are neither peer-reviewed nor mainstreamed. Exactly what proportion of the documented medicinal plant taxa are currently conserved within protected areas is unknown, and baseline monitoring should be promoted to facilitate conservation planning.

Since certain factors (e.g. plant parts removed, harvesting intensity, inherent susceptibility to wound infection, regeneration rates) render some taxa more vulnerable than others to harvesting for medicinal purposes, the IUCN Red List system provides an excellent tool in South Africa for prioritising taxa that are threatened with extinction in the short and medium term. Currently, based on the RLI, less than 2.6% of the medicinal species in South Africa (i.e. <0.3% of the total national indigenous flora) are predicted to be at risk of extinction if no remedial and/or preventative conservation actions are taken. Hence, while medicinal plant

harvesting poses threats to the persistence of socio-economically valuable species, such harvesting is not a significant driver of plant population decline nationally within the context of all (combined) factors threatening South Africa's flora. Readers are referred to Raimondo et al. (2009a) for information on the diversity and scale of such threats. It is important to note, however, that species undergoing decline as a result of unsustainable exploitation to a degree not measureable by the Red List Index (e.g. LC species) may have to undergo substantial changes in population size before their threat status changes sufficiently to reveal the degree of risk to their persistence (Butchart, 2008). Hence, the category of 'Declining' employed in South Africa is a useful means of highlighting which medicinal species are in need of conservation monitoring and intervention. In theory, such action could prevent population declines, ensuring that taxa statuses are not upgraded to threatened or Near Threatened. It thus serves as an 'early warning system' that can inform conservation authorities of which taxa to target.

The harvesting and trade of plant (and animal) material from wild populations for medicinal purposes have been, and remain, a controversial issue, particularly with regard to biodiversity conservation. Socio-economic gains in the short term are often put before long term sustainability of both resources and traditional medicinal practices (Crouch and Smith, 2011). It is our intention that this list of threatened medicinal plant species highlights the importance of conservation management with respect to particular taxa (Appendix A), informing policy development, national and provincial legislation (e.g. TOPS), law enforcement, and most importantly, proactive alternative supply of market-demanded materials through cultivation (Gerstner, 1946; Crouch, 2001; Crouch and Edwards, 2004). Ongoing monitoring of harvesting and trade is required, using techniques comparable to those used historically to quantify these elements, and so allow for the elucidation of trends. Such research should further refine this list and thus re-prioritise species for crucial conservation action.

Appendix A

The medicinal plant species in South Africa that are threatened or are of conservation concern (correct as of September 2012; footnotes indicate status changes since 2009).

Name ^a	Status	Criteria	Family	Recorded in trade
<i>Encephalartos nubimontanus</i>	EW		Zamiaceae	
<i>Encephalartos woodii</i>	EW		Zamiaceae	
<i>Adenium swazicum</i>	CR	A4acd	Apocynaceae	Yes
<i>Agathosma gonaquensis</i>	CR	B1ab(ii,iii,iv,v)	Rutaceae	No
<i>Brackenridgea zanguebarica</i>	CR	A2ad; B1ab(ii,v)	Ochnaceae	Yes
<i>Dioscorea strydomiana</i>	CR	B1ab(v) + 2ab(v); C1	Dioscoreaceae	Yes
<i>Encephalartos cupidus</i>	CR	A2acd; B1ab(ii,iv,v) + 2ab(ii,iv,v)	Zamiaceae	No
<i>Encephalartos dolomiticus</i>	CR	A2d; C1	Zamiaceae	Yes
<i>Encephalartos heenanii</i>	CR	B1ab(ii,iv,v) + 2ab(ii,iv,v)	Zamiaceae	Yes
<i>Encephalartos hirsutus</i>	CR	A4acd; B2ab(iii,iv,v); C1	Zamiaceae	No
<i>Encephalartos laevifolius</i>	CR	A2acde	Zamiaceae	Yes
<i>Encephalartos latifrons</i>	CR	A2ad; B2ab(ii,iii,v); C1 + 2a(i)	Zamiaceae	No
<i>Satyrium rhodanthum</i>	CR	B1ab(i,ii,iii)	Orchidaceae	No
<i>Siphonochilus aethiopicus</i>	CR	A4acd	Zingiberaceae	Yes
<i>Widdringtonia cedarbergensis</i>	CR	A2ab	Cupressaceae	No
<i>Adenia wilmsii</i>	EN	D	Passifloraceae	Yes
<i>Adromischus mammillaris</i>	EN	B1ab(iii) + 2ab(iii)	Crassulaceae	No
<i>Albizia suluensis</i>	EN	B1ab(ii,iii,v) + 2ab(ii,iii,v); C2a(ii)	Fabaceae	Yes
<i>Begonia dregei</i>	EN	C2a(i)	Begoniaceae	Yes
<i>Begonia homonyma</i>	EN	C2a(i)	Begoniaceae	Yes
<i>Cassipourea flanagani</i>	EN	A4acd; C1 + 2a(i)	Rhizophoraceae	Yes
<i>Cyrtanthus suaveolens</i>	EN	B1ab(i,ii,iii)	Amarylloidaceae	No
<i>Encephalartos eugene-maraisii</i>	EN	A2d; C1	Zamiaceae	No
<i>Encephalartos horridus</i>	EN	A2cd	Zamiaceae	No
<i>Encephalartos lebomboensis</i>	EN	A2acd; B1ab(ii,iii,iv,v) + 2ab(ii,iii,iv,v)	Zamiaceae	Yes
<i>Euphorbia woodii</i>	EN	A4cd	Euphorbiaceae	Yes
<i>Haworthia attenuata</i> var. <i>attenuata</i>	EN	A4acd	Asphodelaceae	Yes
<i>Haworthia koelmaniorum</i> var. <i>macmurtryi</i>	EN	B1ab(iii,v) + 2ab(iii,v)	Asphodelaceae	Yes
<i>Ledebouria galpinii</i>	EN	B1ab(iii) + 2ab(iii)	Hyacinthaceae	Yes
<i>Leucospermum conocarpodendron</i> subsp. <i>conocarpodendron</i>	EN	B1ab(i,ii,iii,iv,v) + 2ab(i,ii,iii,iv,v)	Proteaceae	No
<i>Mondia whitei</i>	EN	A2ad	Apocynaceae	Yes
<i>Ocotea bullata</i>	EN	A2bd	Lauraceae	Yes

While it is possible to use market research as a proxy for assessing the condition of medicinal resources in the wild, our current quantitative knowledge of standing stocks is highly deficient both within and outside of theoretically protected areas. For selected species that are currently at high risk, extensive field survey work is needed. In this respect it is necessary to assess and monitor populations (e.g. establishing localities, number of subpopulations, number of mature individuals, genetic diversity, habitat quality, susceptibility to climate change, threats beyond harvesting), and to undertake the autecological research (e.g. generation time, seed viability, pollination ecology, susceptibility to disease post harvesting) that underpins data generation necessary for accurate Red Listing, but which are often difficult to estimate. Management plans for susceptible species are informed by Red Lists, which in turn require quantitative consideration to establish their value limits. The current contribution has thus considered the status, trends, and assessment challenges associated with Red Listing of medicinal plants in South Africa in support of Target 3 of the GSPC (information, research and associated outputs, and methods necessary to implement the Strategy developed and shared) (GPCC, 2010).

Acknowledgements

We thank Tilla Raimondo and Lize von Standen from SANBI's Threatened Species Program for funding and support, and the people who contributed information towards the medicinal plant assessments, particularly the participants of the Durban workshop in January 2008 namely: Tony Cunningham, Mervyn Lötter, Mkhhipheni Ngwenya, Lorraine Mills, Vanessa Brueton and the late Rob Scott-Shaw. Tony Dold is also thanked for information. The reviewers are thanked for constructive comments. VW thanks the University of the Witwatersrand for financial assistance.

Appendix A (continued)

Name ^a	Status	Criteria	Family	Recorded in trade
<i>Sparaxis grandiflora</i> subsp. <i>grandiflora</i>	EN	B1ab(ii,iii,v)	Iridaceae	No
<i>Warburgia salutaris</i>	EN	A2acd	Canellaceae	Yes
<i>Alepidea amatymbica</i> vars	VU	A2d	Apiaceae	Yes
<i>Alepidea macowani</i>	VU	A2ad; B1ab(v)	Apiaceae	Yes
<i>Aloe dichotoma</i>	VU	A3ce	Asphodelaceae	No
<i>Aloe pruinosa</i>	VU	B1ab(ii,iii,v)	Asphodelaceae	Yes
<i>Arctopus dregei</i>	VU	B1ab(i,ii,iii,iv,v)	Apiaceae	No
<i>Argyrolobium longifolium</i>	VU	B1ab(i,iii,v)	Fabaceae	No
<i>Bowiea volubilis</i> subsp. <i>volubilis</i>	VU	A2ad	Hyacinthaceae	Yes
<i>Cassipourea gummiflua</i> var. <i>verticillata</i>	VU	A4acd	Rhizophoraceae	Yes
<i>Ceropegia cimiciodora</i>	VU	B2ab(ii,iii,v)	Apocynaceae	No
<i>Clivia gardenii</i>	VU	A2abcd; B1ab(ii,iv,v)	Amaryllidaceae	Yes
<i>Clivia miniata</i> var. <i>miniata</i>	VU	A2abcd	Amaryllidaceae	Yes
<i>Clivia nobilis</i>	VU	A2cd	Amaryllidaceae	Yes
<i>Clivia robusta</i>	VU	A2cd; B1ab(ii,iii,v)	Amaryllidaceae	Yes
<i>Crinum moorei</i>	VU	A4de	Amaryllidaceae	Yes
<i>Cryptocarya myrtifolia</i>	VU	A2cd	Lauraceae	Yes
<i>Cucumis humifructus</i>	VU	B1ab(ii,v)	Cucurbitaceae	No
<i>Diaphanathe millarii</i>	VU	B1ab(iii,v)	Orchidaceae	No
<i>Dioscorea brownii</i>	VU	D2	Dioscoreaceae	No
<i>Dioscorea sylvatica</i> vars	VU	A2cd	Dioscoreaceae	Yes
<i>Dracosciadium italaie</i>	VU	B1ab(i,ii,iii)	Apiaceae	No
<i>Drimia cooperi</i>	VU	A2ad; C2a(i)	Hyacinthaceae	Yes
<i>Encephalartos altensteinii</i>	VU	A2acd; C1	Zamiaceae	Yes
<i>Encephalartos ghellinckii</i>	VU	C1	Zamiaceae	Yes
<i>Encephalartos humilis</i>	VU	A2acd; C1	Zamiaceae	Yes
<i>Encephalartos ngoyanus</i>	VU	A4acd; C1	Zamiaceae	Yes
<i>Encephalartos paucidentatus</i>	VU	A2acd; B1ab(v) + 2ab(v); C1	Zamiaceae	No
<i>Encephalartos senticosus</i>	VU	A2ace; C1	Zamiaceae	Yes
<i>Eriospermum bracteatum</i>	VU	D2	Eriospermaceae	Yes
<i>Eucomis vandermerwei</i>	VU	B1ab(i,ii,iii,iv,v) + 2ab(i,ii,iii,iv,v)	Hyacinthaceae	Yes
<i>Euphorbia frankiae</i>	VU	A2cd	Euphorbiaceae	Yes
<i>Gasteria croucheri</i>	VU	A2d	Asphodelaceae	Yes
<i>Gerrardanthus tomentosus</i>	VU	D1 + 2	Cucurbitaceae	Yes
<i>Haworthia koelmaniorum</i> var. <i>koelmaniorum</i>	VU	A2acd; C2a(i)	Asphodelaceae	Yes
<i>Haworthia limifolia</i> vars	VU	A2d	Asphodelaceae	Yes
<i>Hydrostachys polymorpha</i>	VU	D2	Hydrostachyaceae	Yes
<i>Knowltonia bracteata</i>	VU	A2d; B2ab(v)	Ranunculaceae	Yes
<i>Leucadendron procerum</i>	VU	A4c	Proteaceae	No
<i>Ocotea kenyensis</i>	VU	D1	Lauraceae	Yes
<i>Oncosiphon africanum</i>	VU	B1ab(ii,iii,iv,v)	Asteraceae	No
<i>Orbea woodii</i>	VU	B1ab(v)	Apocynaceae	Yes
<i>Pleiospilos bolusii</i>	VU	B1ab(iii,iv)	Mesembryanthemaceae	No
<i>Prunus africana</i>	VU	A4acd; C1 + 2a(i)	Rosaceae	Yes
<i>Rhynchosia vendae</i>	VU	B1ab(i,ii,iii,iv,v) + 2ab(i,ii,iii,iv,v)	Fabaceae	Yes
<i>Sceletium expansum</i>	VU	B1ab(ii,iii,iv,v)	Mesembryanthemaceae	Yes
<i>Schlechterina mitostemmatoides</i>	VU	B1ab(ii,iii,v) + 2ab(ii,iii,v)	Passifloraceae	Yes
<i>Stangeria eriopus</i>	VU	A2acd + 4cd	Stangeriaceae	Yes
<i>Steirodiscus tagetes</i>	VU	B1ab(ii,iii,iv,v)	Asteraceae	No
<i>Umtiza listeriana</i>	VU	B1ab(v)	Fabaceae	Yes
<i>Adenia fruticosa</i> subsp. <i>fruticosa</i>	NT	A2c; B1ab(iii,v) + 2ab(iii,v)	Passifloraceae	Yes
<i>Alberta magna</i>	NT	B2ab(ii,iii,v)	Rubiaceae	Yes
<i>Aloe linearifolia</i>	NT	A2c; B1ab(ii,iii,iv,v)	Asphodelaceae	Yes
<i>Aloe micracantha</i>	NT	B1ab(ii,iii,iv,v)	Asphodelaceae	No
<i>Aloe thraskii</i>	NT	A2c	Asphodelaceae	Yes
<i>Anemone fanninii</i>	NT	A2d	Ranunculaceae	Yes
<i>Asparagus stipulaceus</i>	NT	B1ab(ii,iii,iv,v)	Asparagaceae	No
<i>Clivia caulescens</i>	NT	A3d	Amaryllidaceae	Yes
<i>Combretum mkuzense</i>	NT	B1ab(ii,iii,iv,v)	Combretaceae	No
<i>Cotyledon orbiculata</i> var. <i>flanagani</i>	NT	D2	Crassulaceae	Yes
<i>Crinum campanulatum</i>	NT	B1a	Amaryllidaceae	Yes
<i>Curtisia dentata</i>	NT	A2d	Cornaceae	Yes
<i>Cyrtanthus mackenii</i> subsp. <i>cooperi</i>	NT	A2c	Amaryllidaceae	Yes
<i>Drimia sanguinea</i>	NT	A2d	Hyacinthaceae	Yes
<i>Elaeodendron transvaalense</i>	NT	A4ad	Celastraceae	Yes
<i>Encephalartos caffer</i>	NT	A2	Zamiaceae	Yes
<i>Encephalartos ferox</i> ^b	NT	A4d	Zamiaceae	Yes
<i>Encephalartos friderici-guilielmi</i> ^c	NT	A2d	Zamiaceae	Yes
<i>Encephalartos lehmannii</i>	NT	A2d	Zamiaceae	Yes
<i>Encephalartos longifolius</i>	NT	A2d; B1ab(v)	Zamiaceae	Yes
<i>Encephalartos natalensis</i>	NT	A2ad	Zamiaceae	Yes
<i>Erythrophleum lasianthum</i>	NT	A2cd; B1ab(ii,iii,iv,v) + 2ab(ii,iii,iv,v)	Fabaceae	Yes
<i>Eucomis bicolor</i>	NT	A2d	Hyacinthaceae	Yes
<i>Eucomis pallidiflora</i> subsp. <i>pole-evansii</i>	NT	B2ab(v)	Hyacinthaceae	Yes
<i>Euphorbia meloformis</i> subsp. <i>meloformis</i>	NT	B1ab(i,ii,iii,iv,v)	Euphorbiaceae	No

(continued on next page)

Appendix A (continued)

Name ^a	Status	Criteria	Family	Recorded in trade
<i>Gasteria batesiana</i> var. <i>batesiana</i>	NT	A2d; B1 ab(v)	Asphodelaceae	Yes
<i>Gethyllis ciliaris</i> subsp. <i>ciliaris</i>	NT	A2ac; B1ab(ii,iii,iv,v)	Amaryllidaceae	No
<i>Haemanthus deformis</i>	NT	B1ab(v)	Amaryllidaceae	Yes
<i>Haworthia fasciata</i>	NT	B1ab(ii,iii,iv,v)	Asphodelaceae	Yes
<i>Helichrysum cochleariforme</i>	NT	B1ab(ii,iii,iv,v)	Asteraceae	No
<i>Hoodia officinalis</i> subsp. <i>officinalis</i>	NT	B1ab(v)	Apocynaceae	Yes
<i>Hoodia pilifera</i> subsp. <i>pilifera</i>	NT	B1ab(iii,v)	Apocynaceae	Yes
<i>Lithops lesliei</i> subsp. <i>burchellii</i>	NT	D2	Mesembryanthemaceae	Yes
<i>Lithops lesliei</i> subsp. <i>lesliei</i>	NT	A4acd	Mesembryanthemaceae	Yes
<i>Merwillia plumbea</i>	NT	A2bd	Hyacinthaceae	Yes
<i>Pelargonium exhibens</i>	NT	D2	Geraniaceae	Yes
<i>Synaptolepis oliveriana</i>	NT	A4d	Thymelaeaceae	Yes
<i>Urginea lydenburgensis</i>	NT	B1ab(v)	Hyacinthaceae	Yes
<i>Vanilla roscheri</i>	NT	D2	Orchidaceae	No
<i>Acridocarpus natalitius</i> vars	Declining		Malpighiaceae	Yes
<i>Adenia gummifera</i> var. <i>gummifera</i>	Declining		Passifloraceae	Yes
<i>Agathosma betulina</i>	Declining		Rutaceae	Yes
<i>Agathosma crenulata</i>	Declining		Rutaceae	Yes
<i>Aloe cooperi</i> subsp. <i>cooperi</i>	Declining		Asphodelaceae	Yes
<i>Ansellia africana</i>	Declining		Orchidaceae	Yes
<i>Balanites maughamii</i> subsp. <i>maughamii</i>	Declining		Balanitaceae	Yes
<i>Boophone disticha</i>	Declining		Amaryllidaceae	Yes
<i>Callilepis leptophylla</i>	Declining		Asteraceae	Yes
<i>Cassipourea malosana</i>	Declining		Rhizophoraceae	Yes
<i>Crinum bulbispermum</i>	Declining		Amaryllidaceae	Yes
<i>Crinum macowanii</i>	Declining		Amaryllidaceae	Yes
<i>Crinum stuhlmannii</i>	Declining		Amaryllidaceae	Yes
<i>Cryptocarya latifolia</i>	Declining		Lauraceae	Yes
<i>Cryptocarya transvaalensis</i>	Declining		Lauraceae	Yes
<i>Cyathea capensis</i> var. <i>capensis</i>	Declining		Cyathaceae	Yes
<i>Cyrtanthus obliquus</i>	Declining		Amaryllidaceae	Yes
<i>Dioscorea elephantipes</i>	Declining		Dioscoreaceae	Yes
<i>Drimia altissima</i>	Declining		Hyacinthaceae	Yes
<i>Elaeodendron croceum</i>	Declining		Celastraceae	Yes
<i>Eucomis autumnalis</i> subsp.	Declining		Hyacinthaceae	Yes
<i>Eucomis comosa</i> vars	Declining		Hyacinthaceae	Yes
<i>Eucomis montana</i>	Declining		Hyacinthaceae	Yes
<i>Eulophia speciosa</i>	Declining		Orchidaceae	Yes
<i>Euphorbia bupleurifolia</i>	Declining		Euphorbiaceae	Yes
<i>Gunnera perpensa</i>	Declining		Gunneraceae	Yes
<i>Hypoxis hemerocallidea</i>	Declining		Hypoxidaceae	Yes
<i>Ilex mitis</i> var. <i>mitis</i>	Declining		Aquifoliaceae	Yes
<i>Loxostylis alata</i>	Declining		Anacardiaceae	Yes
<i>Newtonia hildebrandtii</i> var. <i>hildebrandtii</i>	Declining		Fabaceae	No
<i>Pronium serratum</i>	Declining		Prioniaceae	Yes
<i>Pterocelastrus rostratus</i>	Declining		Celastraceae	Yes
<i>Rapanea melanophloeos</i>	Declining		Myrsinaceae	Yes
<i>Sandersonia aurantiaca</i>	Declining		Colchicaceae	Yes
<i>Asparagus spinescens</i>	Rare		Asparagaceae	No
<i>Brachystelma meyerianum</i>	Rare		Apocynaceae	No
<i>Cephalaria decurrens</i>	Rare		Dipsacaceae	No
<i>Crassula arborescens</i> subsp. <i>undulatifolia</i>	Critically Rare		Crassulaceae	No
<i>Crassula sarmentosa</i> var. <i>integrifolia</i>	Rare		Crassulaceae	Yes
<i>Disa sanguinea</i>	Rare		Orchidaceae	No
<i>Euphorbia sekukuniensis</i>	Rare		Euphorbiaceae	No
<i>Faurea macnaughtonii</i>	Rare		Proteaceae	Yes
<i>Gasteria bicolor</i> var. <i>liliputana</i>	Rare		Asphodelaceae	No
<i>Greyia flanagani</i>	Rare		Greyiaceae	No
<i>Haemanthus paucifolius</i>	Rare		Amaryllidaceae	No
<i>Huernia pendula</i>	Rare		Apocynaceae	No
<i>Orbea gerstneri</i> subsp. <i>gerstneri</i>	Rare		Apocynaceae	No
<i>Killickia grandiflora</i>	Rare		Lamiaceae	No
<i>Scabiosa transvaalensis</i>	Thr*		Dipsacaceae	No
<i>Tetradenia barberae</i>	Rare		Lamiaceae	No
<i>Thunbergia venosa</i>	Rare		Acanthaceae	Yes
<i>Vitellariopsis dispar</i>	Rare		Sapotaceae	Yes
<i>Acanthosicyos horridus</i> ^d	DDD		Cucurbitaceae	No
<i>Hoodia gordonii</i>	DDD		Apocynaceae	Yes
<i>Isoglossa densa</i>	DDD		Acanthaceae	No
<i>Syringodea flanagani</i>	DDD		Iridaceae	Yes

^aTaxa downgraded to LC: *Acacia mellifera* subsp. *detinens* and *Pelargonium sidoides* (previously Declining); *Orbea paradoxa* and *Tritonia gladiolaris* (previously Rare).

^bPrevious assessment: LC.

^cPrevious assessment: VU C1.

^dPrevious assessment: CR PE.

Appendix B

List of plant families used for traditional medicine in South Africa, the number of species per family, the percentage of the total number of medicinal species used within the family, and the percentage of species threatened and of conservation concern. Proportion of the total medicinal flora in South Africa.

Family	Number of species used per family	% of the total medicinal flora used in South Africa (n = 2062)	% species per family threatened or Near Threatened
Acanthaceae	31	1.5%	
Achariaceae	3	0.1%	
Agapanthaceae	8	0.4%	
Agaricaceae	1	0.05%	
Aizoaceae	4	0.2%	
Alliaceae	10	0.5%	
Amaranthaceae	13	0.6%	
Amaryllidaceae	42	2.0%	26.2%
Anacardiaceae	29	1.4%	
Anemiaceae	1	0.05%	
Annonaceae	8	0.4%	
Anthericaceae	4	0.2%	
Apiaceae	39	1.9%	10.3%
Apocynaceae	104	5.0%	7.7%
Aponogetonaceae	1	0.05%	
Aquifoliaceae	1	0.05%	
Araceae	3	0.1%	
Araliaceae	6	0.3%	
Arecaceae	2	0.1%	
Asparagaceae	24	1.2%	4.2%
Asphodelaceae	83	4.0%	14.5%
Asteraceae	257	12.5%	1.2%
Balanitaceae	2	0.1%	
Balanophoraceae	1	0.05%	
Balsaminaceae	1	0.05%	
Begoniaceae	3	0.1%	66.7%
Behniaceae	1	0.05%	
Bignoniaceae	3	0.1%	
Blechnaceae	1	0.05%	
Boraginaceae	14	0.7%	
Brassicaceae	6	0.3%	
Buddlejaceae	7	0.3%	
Burseraceae	4	0.2%	
Buxaceae	1	0.05%	
Cactaceae	1	0.05%	
Campanulaceae	8	0.4%	
Canellaceae	1	0.05%	100.0%
Capparaceae	19	0.9%	
Caryophyllaceae	15	0.7%	
Celastraceae	20	1.0%	5.0%
Celtidaceae	5	0.2%	
Chenopodiaceae	3	0.1%	
Chrysobalanaceae	2	0.1%	
Clusiaceae	2	0.1%	
Colchicaceae	4	0.2%	
Combretaceae	16	0.8%	6.3%
Commelinaceae	5	0.2%	
Convolvulaceae	17	0.8%	
Cornaceae	1	0.05%	100.0%
Crassulaceae	32	1.6%	6.3%
Cucurbitaceae	24	1.2%	12.5%
Cunoniaceae	1	0.05%	
Cupressaceae	2	0.1%	50.0%
Cyatheaceae	2	0.1%	
Cyperaceae	7	0.3%	
Dennstaedtiaceae	1	0.05%	
Dioscoreaceae	8	0.4%	37.5%
Dipsacaceae	8	0.4%	
Dracaenaceae	4	0.2%	
Dryopteridaceae	3	0.1%	
Ebenaceae	15	0.7%	
Elatinaceae	1	0.05%	
Equisetaceae	1	0.05%	
Erioseptaceae	6	0.3%	16.7%

Appendix B (continued)

Family	Number of species used per family	% of the total medicinal flora used in South Africa (n = 2062)	% species per family threatened or Near Threatened
Euphorbiaceae	65	3.2%	4.6%
Fabaceae	159	7.7%	3.1%
Flacourtiaceae	3	0.1%	
Gentianaceae	10	0.5%	
Geraniaceae	37	1.8%	2.7%
Gesneriaceae	1	0.05%	
Greyiaceae	2	0.1%	
Gunneraceae	1	0.05%	
Haemodoraceae	1	0.05%	
Hernandiaceae	1	0.05%	
Heteropyxidaceae	2	0.1%	
Hyacinthaceae	44	2.1%	20.5%
Hydnoraceae	2	0.1%	
Hydrostachyaceae	1	0.05%	100.0%
Hypericaceae	1	0.05%	
Hypoxidaceae	10	0.5%	
Icacinaceae	5	0.2%	
Iridaceae	41	2.0%	2.4%
Lamiaceae	68	3.3%	
Lauraceae	7	0.3%	42.9%
Lecythidaceae	1	0.05%	
Lentibulariaceae	3	0.1%	
Lessoniaceae	1	0.05%	
Linaceae	2	0.1%	
Lobeliaceae	10	0.5%	
Loranthaceae	2	0.1%	
Lycopodiaceae	4	0.2%	
Lythraceae	1	0.05%	
Maesaceae	2	0.1%	
Malpighiaceae	3	0.1%	
Malvaceae	50	2.4%	
Melastomataceae	2	0.1%	
Meliaceae	6	0.3%	
Meliantaceae	7	0.3%	
Menispermaceae	12	0.6%	
Menyanthaceae	1	0.05%	
Mesembryanthemaceae	22	1.1%	18.2%
Molluginaceae	6	0.3%	
Monimiaceae	1	0.05%	
Moraceae	9	0.4%	
Musaceae	1	0.05%	
Myricaceae	4	0.2%	
Myrothamnaceae	1	0.05%	
Myrsinaceae	2	0.1%	
Myrtaceae	4	0.2%	
Nymphaeaceae	2	0.1%	
Ochnaceae	7	0.3%	14.3%
Olaceae	2	0.1%	
Oleaceae	8	0.4%	
Orchidaceae	63	3.1%	4.8%
Orobanchaceae	11	0.5%	
Oxalidaceae	4	0.2%	
Passifloraceae	6	0.3%	50.0%
Pedaliaceae	7	0.3%	
Phyllanthaceae	9	0.4%	
Phytolaccaceae	2	0.1%	
Piperaceae	1	0.05%	
Pittosporaceae	1	0.05%	
Plantaginaceae	1	0.05%	
Plumbaginaceae	1	0.05%	
Poaceae	25	1.2%	
Podocarpaceae	3	0.1%	
Polygalaceae	21	1.0%	
Polygonaceae	9	0.4%	
Polypodiaceae	2	0.1%	
Portulacaceae	7	0.3%	
Potamogetonaceae	1	0.05%	
Prioniaceae	1	0.05%	
Proteaceae	13	0.6%	15.4%
Ptaeroxylaceae	1	0.05%	
Pteridaceae	9	0.4%	
Putranjivaceae	1	0.05%	
Ranunculaceae	12	0.6%	16.7%

(continued on next page)

Appendix B (continued)

Family	Number of species used per family	% of the total medicinal flora used in South Africa (n = 2062)	% species per family threatened or Near Threatened
Restionaceae	1	0.05%	
Rhamnaceae	8	0.4%	
Rhizophoraceae	6	0.3%	33.3%
Rosaceae	11	0.5%	9.1%
Rubiaceae	46	2.2%	2.2%
Rutaceae	21	1.0%	4.8%
Salicaceae	9	0.4%	
Salvadoraceae	2	0.1%	
Samydaceae	1	0.05%	
Santalaceae	6	0.3%	
Sapindaceae	10	0.5%	
Sapotaceae	9	0.4%	
Scrophulariaceae	24	1.2%	
Selaginellaceae	3	0.1%	
Smilacaceae	1	0.05%	
Solanaceae	14	0.7%	
Stangeriaceae	1	0.05%	100.0%
Strelitziaceae	2	0.1%	
Strychnaceae	7	0.3%	
Tamaricaceae	1	0.05%	
Tecophilaeaceae	1	0.05%	
Theophrastaceae	1	0.05%	
Thymelaeaceae	15	0.7%	13.3%
Typhaceae	1	0.05%	
Urticaceae	5	0.2%	
Valerianaceae	1	0.05%	
Velloziaceae	2	0.1%	
Verbenaceae	5	0.2%	
Violaceae	2	0.1%	
Viscaceae	6	0.3%	
Vitaceae	12	0.6%	
Zamiaceae	24	1.2%	91.7%
Zingiberaceae	1	0.05%	100.0%
Zygophyllaceae	2	0.1%	

References

- Anonymous, c.1994. Table V: Useful plant resources to communities in and around Songimvelo Game Reserve. Unpublished species list.
- Arnold, T.H., Prentice, C.A., Hawker, L.C., Snyman, E.E., Tomalin, M., Crouch, N.R., Pottas-Bircher, C., 2002. Medicinal and magical plants of southern Africa: an annotated checklist. *Strelitzia*, 13. National Botanical Institute, Pretoria.
- Bircher, C., Prentice, C., Crouch, N., Symmonds, R., 1998. Conservation concerns for *Bowiea volubilis*, an unusually succulent member of the Hyacinthaceae. *Herbertia* 53, 81–88.
- Botha, J., Witkowski, E.T.F., Shackleton, C.M., 2001. An inventory of medicinal plants traded on the western boundary of the Kruger National Park, South Africa. *Koedoe* 44 (2), 7–46.
- Botha, J., Witkowski, E.T.F., Shackleton, C.M., 2004a. The impact of commercial harvesting on *Warburgia salutaris* ('pepper-bark tree') in Mpumalanga, South Africa. *Biodiversity and Conservation* 13, 1675–1698.
- Botha, J., Witkowski, E.T.F., Shackleton, C.M., 2004b. Market profiles and trade in medicinal plants in the Lowveld, South Africa. *Environmental Conservation* 31, 38–46.
- Brandt, H.D., Osuch, E., Mathibe, L., Tsipa, P., 1995. Plants associated with accidentally poisoned patients at Ga-Rankuwa hospital, Pretoria. *South African Journal of Science* 91, 57–59.
- Brenan, J.P.M., 1970. Leguminosae. *Flora Zambesiaca* 3 (1).
- Brueton, V.J., in preparation. Trade in commonly used medicinal bulbs: value and ecological implications. MSc Dissertation, University of the Witwatersrand, Johannesburg.
- Bubb, P.J., Butchart, S.H.M., Collen, B., Dublin, H., Kapos, V., Pollock, C., Stuart, S.N., Vié, J.-C., 2009. IUCN Red List Index – Guidance for National and Regional Use. IUCN, Gland, Switzerland (http://intranet.iucn.org/webfiles/doc/SpeciesProg/RLI_Guidelines_Final_4march09.pdf).
- Butchart, S.H.M., 2008. Red List Indices to measure the sustainability of species use and impacts of invasive alien species. *Bird Conservation International* 18, S245–S262.
- Cocks, M.L., 1996. Towards an understanding of *amaYeza esiXhosa* Stores (African Chemists). M.A. thesis. Rhodes University, Grahamstown.
- Cooper, K., 1979. A muti man's stinkwood racket. *African Wildlife* 33, 8.
- Cousins, S.R., Williams, V.L., Witkowski, E.T.F., 2011. Quantifying the trade in cycads (*Encephalartos* species) in the traditional medicine markets of Johannesburg and Durban, South Africa. *Economic Botany* 65, 356–370.
- Cousins, S.R., Williams, V.L., Witkowski, E.T.F., 2012. Uncovering the cycad taxa (*Encephalartos* species) traded for traditional medicine in Johannesburg and Durban, South Africa. *South African Journal of Botany* 78, 129–138.
- Crouch, N.R., 2001. The role of horticulture in threatened species conservation. *FSA ethnomedicinal taxa in focus. SABONET News* 6, 185–188.
- Crouch, N.R., Edwards, T., 2004. Ethnomedicinal (*muthi*) plant nurseries. In: Lawes, M.J., Eeley, H.A.C., Shackleton, C.M., Geach, B.G.S. (Eds.), *Indigenous Forests and Woodlands in South Africa: Policy, People and Practice*. University of KwaZulu-Natal Press, Pietermaritzburg, pp. 658–661 (675–678).
- Crouch, N.R., Hutchings, A., 1999. Zulu healer *muthi* gardens: inspiration for botanic garden displays and community outreach projects. *Proceedings of the 5th International Botanic Gardens Conservation Congress* (http://www.bgci.org/congress/congress_1998_cape/html/crouch.htm).
- Crouch, N.R., Krynauw, S., 1999. The high altitude pineapple lily *Eucomis vandermerwei*, a rare endemic from South Africa. *Herbertia* 54, 133–138.
- Crouch, N.R., Smith, G.F., 2011. Informing and influencing the interface between biodiversity science and biodiversity policy in South Africa. *Botanical Journal of the Linnean Society* 16 (6), 301–309.
- Crouch, N.R., Lötter, M., Krynauw, S., Pottas-Bircher, C., 2000. *Siphonochilus aethiopicus* (Zingiberaceae), the prized *indungulu* of the Zulu – an overview. *Herbertia* 55 (89), 115–129.
- Crouch, N.R., Mulholland, D.A., Pohl, T.L., Ndlovu, E., 2003a. The ethnobotany and chemistry of the genus *Clivia* (Amaryllidaceae). *South African Journal of Botany* 69 (2), 144–147.
- Crouch, N.R., Smith, G.F., Lötter, M., Symmonds, R., 2003b. *Encephalartos woodii*: the first historically documented *muthi* casualty in southern Africa. *Encephalartos* 73 (22), 25–28.
- Crouch, N.R., Langlois, A., Mulholland, D.A., Nair, J.J., 2005a. A novel alkylamide from the leaves of *Acmella caulirhiza* (Asteraceae), a traditional surface analgesic. *South African Journal of Botany* 71, 228–230.
- Crouch, N.R., Pohl, T.L., Mulholland, D.A., Ndlovu, E., 2005b. Alkaloids from three ethnomedicinal *Haemanthus* species: *H. albiflos*, *H. deformis* and *H. pauciflorus* (Amaryllidaceae). *South African Journal of Botany* 71, 49–52.
- Cunningham, A.B., 1988. An investigation of the herbal medicine trade in Natal/KwaZulu. *Investigational Report*, vol. 29. Institute of Natural Resources, Pietermaritzburg.
- Dally, K., 1984. Illegal debarking of stinkwood. *Forestry News* 2/85, 23.
- Dicke, B.H., 1926. A Bavenda sacred object. *South African Journal of Science* 22, 935–936.
- Dold, A.P., Cocks, M.L., 2000. The medicinal use of some weeds, problem and alien plants in the Grahamstown and Peddie districts of the Eastern Cape, South Africa. *South African Journal of Science* 96, 467–473.
- Dold, A.P., Cocks, M.L., 2001. Traditional veterinary medicine in the Alice district of the Eastern Cape Province, South Africa. *South African Journal of Science* 97, 375–379.
- Dold, A.P., Cocks, M.L., 2002. The trade in medicinal plants in the Eastern Cape Province, South Africa. *South African Journal of Science* 98, 589–597.
- Douwes, E., Crouch, N.R., Edwards, T.J., Mulholland, D.A., 2008. Regression analyses of southern African ethnomedicinal plants: informing the targeted selection of bioprospecting and pharmacological screening subjects. *Journal of Ethnopharmacology* 119, 356–364.
- Dzerefos, C.M., Witkowski, E.T.F., 2001. Density and potential utilization of medicinal grassland plants from Abe Bailey Nature Reserve, South Africa. *Biodiversity and Conservation* 10, 1875–1896.
- Fairbanks, D.H.K., Thompson, M.W., Vink, D.E., Newby, T.S., Van den Berg, H.M., Everard, D.A., 2000. The South African Land-Cover Database: a synopsis of the landscape. *South African Journal of Science* 96, 69–82.
- Fourcade, H.G., 1889. Report on the Natal Forests. Natal Blue Book. Watson, Government Printer, Pietermaritzburg.
- Geldenhuys, C.J., 2004. Meeting the demand for *Ocotea bullata* bark. Implications for the conservation of high-value and medicinal tree species. In: Lawes, M.J., Eeley, H.A.C., Shackleton, C.M., Geach, B.G.S. (Eds.), *Forests and Woodlands in South Africa: Policy, People and Practice*. University of Natal Press, Pietermaritzburg, pp. 517–550.
- Gerstner, J., 1938. A preliminary check list of Zulu names of plants. *Bantu Studies* 12, 215–236.
- Gerstner, J., 1946. Some factors affecting the perpetuation of our indigenous silva. *Journal of the South African Forestry Association* 13, 4–11.
- Global Partnership for Plant Conservation (GPPC), 2010. The Global Strategy for Plant Conservation (GSPC). <http://www.plants2020.net/gspc-targets/> (accessed 20 September).
- Hanekom, C., 1967. Tradisionele geneeskunde by enkele Noord-Sothostamme. *African Studies* 26, 37–42.
- Hutchings, A., Scott, A.H., Lewis, G., Cunningham, A.B., 1996. *Zulu Medicinal Plants: an Inventory*. University of Natal Press, Pietermaritzburg.
- Kamatou, G.P.P., Makunga, N.P., Ramogola, W.P.N., Viljoen, A.M., 2008. South African *Salvia* species: a review of biological activities and phytochemistry. *Journal of Ethnopharmacology* 119, 664–672.
- Keith, M., Warren, M., 2007. Can anthropogenic variables be used as threat proxies for South African plant richness? *Bothalia* 37, 79–88.
- King, N.L., 1939. The Knysna forests and the woodcutter problem. *Journal of the South African Forestry Association* 3, 6–15.
- King, N.L., 1941. The exploitation of the indigenous forests of South Africa. *Journal of the South African Forestry Association* 6, 23–48.
- Kroon, N.V., 1999. A checklist of the flora of Grid 2627DD, Sasolburg, Free State: a springboard for conservation at Wonderwater strip mine, Sasol Coal. *Plantlife Supplementary* 1, 1–36.
- Lourens, A.C.U., Viljoen, A.M., Van Heerden, F.R., 2008. South African *Helichrysum* species: a review of the traditional uses, biological activity and phytochemistry. *Journal of Ethnopharmacology* 119, 630–652.
- Loxton, Venn and Associates, 1994. Driekoppies Dam Valuation and Resettlement Programme. Traditional and Medicinal Plants Mitigation Options. Report to Department of Home Affairs, Kangwane.

- Lubbe, W.A., Mostert, G.P., 1991. Rate of *Ocotea bullata* decline in association with *Phytophthora cinnamomi* at three study sites in the southern Cape indigenous forests. *South African Forestry Journal* 159, 17–24.
- Makgagga, C., 1995. Useful plants of the Northern Province: a preliminary list of North Sotho plant names and their uses. *Plantlife* 13, 27–29.
- Makunga, N.P., Philander, L.E., Smith, M., 2008. Current perspectives on an emerging formal natural products sector in South Africa. *Journal of Ethnopharmacology* 119, 365–375.
- Mander, M., 1997. Medicinal Plant Marketing and Strategies for Sustaining the Plant Supply in the Bushbuckridge Area and Mpumalanga Province, South Africa. Department of Water Affairs and Forestry, Pretoria.
- Mander, M., 1998. Marketing of Indigenous Medicinal Plants in South Africa: a Case Study in KwaZulu-Natal. Food and Agriculture Organization of the United Nations, Rome.
- Mander, M., Ntuli, L., Diederichs, N., Mavundla, K., 2007. Economics of the traditional medicine trade in South Africa. In: Harrison, S., Bhana, R., Ntuli, A. (Eds.), *South African Health Review 2007*. Health Systems Trust, Durban (http://www.hst.org.za/uploads/files/chap13_07.pdf).
- Matsiliza, B., 1997. Plants Used in Traditional Medicine in Grahamstown. B.Sc. Honours Dissertation, Rhodes University, Grahamstown.
- McCracken, D.P., 1986. The indigenous forests of colonial Natal and Zululand. *Natalia* 16, 19–38.
- McCraw, L.J., Lall, N., Meyer, J.J.M., Eloff, J.N., 2008. The potential of South African plants against *Mycobacterium* infections. *Journal of Ethnobiology* 482–500.
- Moeng, E.T., Potgieter, M.J., 2011. The trade of medicinal plants by muthi shops and street vendors in Limpopo Province, South Africa. *Journal of Medicinal Plant Research* 5 (4), 558–564.
- More, G., Tshikalange, T.E., Lall, N., Botha, F., Meyer, J.J.M., 2008. Antimicrobial activity of medicinal plants against microorganisms. *Journal of Ethnopharmacology* 119, 473–477.
- Morty, K.E., Johnson, D.N., 1987. The status and conservation of the black stinkwood in Natal. *Lammerger* 38, 60–67.
- Mucina, L., Rutherford, M.C. (Eds.), 2006. The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia*, 19. SANBI, Pretoria.
- Netshingani, E.N., 1981. Notes on the uses of indigenous trees in Venda. *Journal of Dendrology* 1, 12–17.
- Newton, D., 1993. Visit to Blouberg on a collecting trip with healers. Unpublished notes.
- Oatley, T.B., 1984. Exploitation of a new niche by the Rameron Pigeon *Columba arquatrix* in Natal. *Proceedings of the Fifth Pan-African Ornithological Congress*. Southern African Ornithological Society, Johannesburg, pp. 323–330.
- Palmer, E., Pitman, N., 1972. *Trees of Southern Africa*. A.A. Balkema, Cape Town.
- Phillips, J.F.V., 1931. Forest-succession and ecology in the Knysna region. *Memoirs of the Botanical Survey of South Africa*, 14. Department of Agriculture, The Government Printer, Pretoria.
- Pooley, E., 1993. *The Complete Field Guide to Trees of Natal, Zululand and Transkei*. Natal Flora Publications Trust, Durban.
- Pooley, E., 1998. *A Field Guide to Wild Flowers of KwaZulu-Natal and the Eastern Region*. Natal Flora Publications Trust, Durban.
- Raimondo, D.C., Donaldson, J.S., 2003. Responses of cycads with different life histories to the impact of plant collecting: stimulation models to determine important life history stages and population recovery times. *Biological Conservation* 111, 345–358.
- Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), 2009a. Red List of South African Plants. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria.
- Raimondo, D., Victor, J.E., Scott-Shaw, C.R., Lötter, M., Dold, A.P., Pfaf, M.F., Burrows, J.E., 2009b. *Bowiea volubilis* subsp. *volubilis*. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria.
- Rivers, M.C., Taylor, L., Brummitt, N.A., Meagher, T.R., Roberts, D.L., Lughadha, E.N., 2011. How many herbarium specimens are needed to detect threatened species? *Biological Conservation* 144, 2541–2547.
- Rycroft, H.B., 1944. The Karkloof Forest, Natal. *Journal of the South African Forestry Association* 11, 13–25.
- SANBI, 2012. Red List of South African plants version 2012.1. Downloaded from <http://redlist.sanbi.org> (on 2012/09/05).
- Schmidt, E., Lötter, M., McClelland, W., 2002. *Trees and Shrubs of Mpumalanga and Kruger National Park*. Jacana, Johannesburg.
- Scott-Shaw, C.R., 1999. *Rare and Threatened Plants of KwaZulu-Natal and Neighbouring Regions*. KwaZulu-Natal Nature Conservation Service, Pietermaritzburg.
- Sim, T.R., 1906. The Forests and Flora of the Colony of the Cape of Good Hope. Government of the Cape of Good Hope, Pietermaritzburg.
- Smith, G.F., Crouch, N.R., 1995. Notes on *Aloe pruinosus* Reyn. (Aloaceae), a rare and little-known maculate from the midlands of KwaZulu-Natal, South Africa. *Aloe* 32, 66–69.
- Sobiecki, J.F., 2002. A preliminary inventory of plants used for psychoactive purposes in southern African healing traditions. *Transactions of the Royal Society of South Africa* 57, 1–24.
- Statistics South Africa, 2004. *South African Statistics, 2004/05*. P0302 www.statssa.gov.za/Publications.
- Statistics South Africa, 2007. *Community Survey 2007*. CS2007Basic www.statssa.gov.za/Publications.
- Taylor, H.C., 1961. The Karkloof forest: a plea for its protection. *Forestry in South Africa* 1, 123–134.
- Taylor, H.C., 1963. A report on the Nxamalala Forest. *Forestry in South Africa* 2, 29–43.
- Te Beest, M., 2003. The impact of medicinal plant use on biodiversity: a case study in Hluhluwe-Umfolozi Park, KwaZulu-Natal, South Africa. Unpublished report, KZN Wildlife, Pietermaritzburg.
- Todd, C.B., 1999. Mutavhatsindi. A magical medicinal tree from the Soutpansberg. *Veld & Flora* 85 (1), 17.
- Todd, C.B., Khorombbi, K., Van der Waal, B.C., Weisser, P.J., 2004. Conservation of woodland and biodiversity. A complementary traditional and western approach towards protecting *Brackenridgea zanguebarica*. In: Lawes, M.J., Eeley, H.A.C., Shackleton, C.M., Geach, B.G.S. (Eds.), *Indigenous Forests and Woodlands in South Africa: Policy, People and Practice*. University of KwaZulu-Natal Press, Pietermaritzburg, pp. 737–750.
- Van Jaarsveld, E., 1992. *Bowiea gariepensis* and *B. volubilis*. *British Cactus and Succulent Journal* 10 (4), 97.
- Van Vuuren, S.F., 2008. Antimicrobial activity of South African medicinal plants. *Journal of Ethnopharmacology* 119, 462–472.
- Van Wyk, B.-E., 2008. A review of Khoi-San and Cape Dutch medical ethnobotany. *Journal of Ethnopharmacology* 119, 331–341.
- Van Wyk, B.-E., Van Oudtshoorn, B., Gericke, N., 1997. *Medicinal Plants of South Africa*. Briza, Pretoria.
- Veale, D.J.H., Furman, K.I., Oliver, D.W., 1992. South African traditional herbal medicines used during pregnancy and childbirth. *Journal of Ethnopharmacology* 36, 185–191.
- Verschaeve, L., Van Staden, J., 2008. Mutagenic and antimutagenic properties of extracts from South African traditional medicinal plants. *Journal of Ethnopharmacology* 575–587.
- Victor, J.E., 2002. South Africa. In: Golding, J.S. (Ed.), *Southern African Red Data Lists: SABONET Report*, 14, pp. 93–120.
- Victor, J.E., Keith, M.A., 2004. The Orange List: a safety net for biodiversity in South Africa. *South African Journal of Science* 100, 139–141.
- Von Ahlefeldt, D., Crouch, N.R., Nichols, G., Symmonds, R., McKean, S., Sibiya, H., Cele, M.P., 2003. *Medicinal Plants Traded on South Africa's Eastern Seaboard*. Ethekwini Parks Department and University of Natal, Durban, p. 267.
- Von Staden, L., Raimondo, D., Foden, W., 2009a. Introduction. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria, pp. 1–5.
- Von Staden, L., Raimondo, D., Foden, W., 2009b. Assessing the conservation status of South Africa's medicinal plant taxa. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria, p. 18.
- Von Staden, L., Raimondo, D., Foden, W., 2009c. Approach to Red List assessments. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria, pp. 6–16.
- Watt, J.M., Breyer-Brandwijk, M.G., 1962. *The Medicinal and Poisonous Plants of Southern and Eastern Africa*. E. and S. Livingstone Ltd., Edinburgh.
- Williams, V.L., 2003. *Hawkers of health: an investigation of the Faraday Street Traditional Medicine Market in Johannesburg*. Gauteng. Plant Ecology and Conservation Series No. 15. Unpublished report to the Gauteng Directorate for Nature Conservation, DACEL, Johannesburg.
- Williams, V.L., 2005. *Clivia* under threat. *Clivia* 7, 12–16.
- Williams, V.L., 2007. The design of a risk assessment model to determine the impact of the herbal medicine trade on the Witwatersrand on resources of indigenous plant species. PhD Thesis. University of the Witwatersrand, Johannesburg. <http://wiredspace.wits.ac.za/handle/10539/5313>.
- Williams, V.L., 2009. *Elephantorrhiza elephantina*. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria.
- Williams, V.L., 2010. Threatened ethnomedicinal plants of South Africa: Red-Listed species, conservation 'hot spots' and the impact of harvesting. Abstract from the 2nd International Symposium in Medicinal Plants, Their Cultivation and Aspects of Use. 3–4 November 2010. Petra, Jordan.
- Williams, V.L., Geldenhuys, C.J., 2004. Preliminary results of bark recovery patterns for 5 species, 6 years after bark removal. Unpublished report, FRP-DFID R8305 Bark study.
- Williams, V.L., Raimondo, D., 2009. *Brackenridgea zanguebarica*. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria.
- Williams, V.L., Balkwill, K., Witkowski, E.T.F., 2000. Unravelling the commercial market for medicinal plants and plant parts on the Witwatersrand, South Africa. *Economic Botany* 54, 310–327.
- Williams, V.L., Balkwill, K., Witkowski, E.T.F., 2001. A lexicon of plants traded in the Witwatersrand *umuthi* shops. *Bothalia* 31, 71–98.
- Williams, V.L., Balkwill, K., Witkowski, E.T.F., 2007a. Size-class prevalence of bulbous and perennial herbs sold in the Johannesburg medicinal plant markets between 1995 and 2001. *South African Journal of Botany* 73, 144–145.
- Williams, V.L., Witkowski, E.T.F., Balkwill, K., 2007b. Relationship between bark thickness and diameter at breast height for six tree species used medicinally in South Africa. *South African Journal of Botany* 73, 449–465.
- Williams, V.L., Raimondo, D., Crouch, N.R., Cunningham, A.B., Scott-Shaw, C.R., Lötter, M., Ngwenya, A.M., Dold, A.P., 2009. *Ocotea bullata*. In: Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C., Kamundi, D.A., Manyama, P.A. (Eds.), *Red List of South African Plants*. *Strelitzia*, 25. South African National Biodiversity Institute, Pretoria.