

## Research Article

**Cite this article:** Chetty D, Datta A, Kumschick S, Wilson JRU, Nchu F, and Geerts S (2024) Regulatory options for cultivars and hybrids of invasive plant species—the South African experience. *Invasive Plant Sci. Manag* **17**: 228–238. doi: [10.1017/inp.2024.24](https://doi.org/10.1017/inp.2024.24)

Received: 31 May 2024

Accepted: 6 August 2024

First published online: 29 October 2024

### Associate Editor:

Jacob N. Barney, Virginia Tech







### Keywords:

Invasive taxa; regulatory approaches; regulatory lists; stakeholder negotiations; sterile cultivars and hybrids

### Corresponding author:

Duran Chetty; Email: [duranchetty5@gmail.com](mailto:duranchetty5@gmail.com)

# Regulatory options for cultivars and hybrids of invasive plant species—the South African experience

Duran Chetty<sup>1</sup> , Arunava Datta<sup>2</sup> , Sabrina Kumschick<sup>3</sup> , John R.U. Wilson<sup>4</sup> , Felix Nchu<sup>5</sup>  and Sjirk Geerts<sup>6</sup> 

<sup>1</sup>Doctoral Candidate, Department of Horticultural Sciences, Cape Peninsula University of Technology, Bellville Campus, Cape Town, South Africa; and South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa; <sup>2</sup>Researcher and Assistant Professor, South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa; Department of Botany, Raja Narendralal Khan Women's College, Gope Palace, Vidyasagar University, Paschim Medinipur, India; and Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa; <sup>3</sup>Senior Researcher, Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa; and South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town, South Africa; <sup>4</sup>Principal Scientist and Extraordinary Professor, South African National Biodiversity Institute, Kirstenbosch Research Centre, Cape Town; and Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa; <sup>5</sup>Associate Professor, Department of Horticultural Sciences, Cape Peninsula University of Technology, Bellville Campus, Cape Town, South Africa and <sup>6</sup>Associate Professor, Department of Conservation and Marine Sciences, Cape Peninsula University of Technology, Cape Town, South Africa

## Abstract

Invasive plant taxa are generally regulated at the species level, without considering infra- or interspecific variation. However, cultivars or hybrids can pose a lower risk of invasion, for example, due to sterility. We evaluate six general approaches to regulating cultivars and hybrids: (1) Globally Guilty by Association; (2) Nationally Guilty by Association; (3) Guilty until Proven Innocent; (4) Negotiated Guilt; (5) Claimed to be Innocent; and (6) Innocent until Proven Guilty. We discuss these approaches in the context of South Africa (which has a typified Negotiated Guilt approach). Following negotiations since 2001 between the South African horticultural industry/green industry and legislators, an unofficial consensus list of “presumed sterile” cultivars and hybrids was produced in 2014 containing 187 entities from 34 taxa. In 2020, this was reduced to 157 entities from 16 taxa. But the evidence supporting the original lists and the subsequent revisions was not published. To address this issue, we developed a generic pro forma (template) for reporting sterility based on observations and/or experiments on: flowering, fruiting, pollen, and seeds; the potential for vegetative propagation; and the potential for genetic changes (including hybridization and reversion to fertility). We recommend that such information be incorporated into risk analyses conducted specifically for infra- and inter-specific entities, and only if the risk of a harmful invasion is demonstrated to be acceptably low or can be easily mitigated should such entities be exempted from regulation. This will be time-consuming, but, by setting out the evidence clearly, the approach is transparent and provides a clear route for stakeholders to seek exemptions for entities of importance. In conclusion, although we suspect the simplicity of the Negotiated Guilt approach is desirable to many stakeholders, and is the approach currently adopted in South Africa, we recommend a shift toward the Guilty until Proven Innocent approach.

## Introduction

Biological invasions have negative impacts on ecosystems and economies (IPBES 2023; Pyšek et al. 2012; Vilà et al. 2011). To combat or prevent these negative effects, various regulations at different scales (local, regional, or national) have been developed (Hulme et al. 2018; Turbelin et al., 2017), often including the use of regulatory species lists (García-de-Lomas and Vilà 2015; Pergl et al. 2016). These regulatory lists usually focus on taxa that are known or perceived to be harmful (cf. Kumschick et al. [2024] for a discussion on lists of taxa that are of low risk), although all such lists can be complex to develop and will have uncertainties (McGeoch et al. 2012). One particular issue is that non-native plant taxa are assessed for their invasion risk at the species level (Kumschick and Richardson 2013) and thus are also regulated at this level; that is, species are listed, with minor attention given to infraspecific (such as cultivars, forms, or varieties) or interspecific (hybrids) entities (e.g., in Poland [Tokarska-Guzik et al. 2021] and Japan [Mizutani and Goka 2010]).

At the population level, many plant species are individually clustered into distinct genetic lineages across their geographic ranges, suggesting adaptation to local conditions (Hereford 2009;

© The Author(s), 2024. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



### Management Implications

We outline six approaches to regulating cultivars and/or hybrids of invasive plant species. Each approach requires different levels of resources and evidence and results in different levels of risk. We outline the consequences of the different approaches so regulators and stakeholders can choose the best option for their needs.

In general, we recommend a Guilty until Proven Innocent approach. All cultivars or hybrids that are related to at least one regulated taxon should be similarly regulated unless and until there is documented evidence that the cultivar/hybrid is not invasive (i.e., risk analyses should be conducted for each exempted entity). This approach aims to restrict harmful invasions but also to provide stakeholders with a process to advocate for the exemption of entities that are valuable to them (albeit one that requires substantial evidence for a change to be made). We provide a pro forma (template) to support reporting low levels of invasiveness based on observations and/or experiments. We note that, to be effective, the approach requires that the public (and nursery customers) are aware of the process and that sterile cultivars/hybrids can be easily distinguished in practice.

In South Africa, exemptions have been made for several cultivars and hybrids on the basis of presumed sterility. These exemptions emerged from a series of lengthy negotiations between the South African horticulture industry and the regulators and can be typified as a Negotiated Guilt approach. This approach requires fewer resources and is much simpler to execute, but likely leads to more entities being considered safe that are in fact harmful. We motivate for a shift to the Guilty until Proven Innocent approach.

Leimu and Fischer 2008; Linhart and Grant 1996). Gotelli and Stanton-Geddes (2015) suggest that infraspecific variation needs to be considered when modeling shifts in the geographic ranges of plant populations. There can be large variation among infra- and inter specific entities and the parent species. For example, infraspecific entities of port jackson [*Acacia saligna* (Labill.) H.L.Wendl.] occupy different bioclimatic niches within the species' native range, and there is evidence that this is also the case in its invasive range (Thompson et al. 2011). Infraspecific variation can also translate to differences in the impacts caused by invasive taxa, such as spineless cultivars of mission prickly pear [*Opuntia ficus-indica* (L.) Mill.] which are presumed to be non-invasive due to increased herbivory that regulates these populations (Novoa et al. 2018; Zimmermann and Granata 2002).

Hybridization in plants can occur between (interspecific) or within (intraspecific) species, resulting in several possible genetic changes (Landry et al. 2007) that can increase or reduce fitness (Charlesworth 2009). Such genetic changes can influence invasion success (Buhk and Thielsch 2015), and there are many examples of invasive taxa that evolved after intertaxon hybridization (Dlugosch et al. 2015; Ellstrand and Schierenbeck 2000; Hovick and Whitney 2014). Infraspecific hybridization can also promote invasiveness, as seen in the case of calley pear (*Pyrus calleryana* Dence.) (Culley and Hardiman 2009). Conversely, hybridization has been used as a tool to develop sterile hybrids of known invasive taxa, such as the sterile hybrids of Britton's wild petunia (*Ruellia simplex* C. Wright), which were fruitless with low pollen viability (Freyre et al. 2012). It is evident that infra- and interspecific entities can either pose a lower or higher invasion risk. Therefore, not considering infra- or interspecific variation when developing invasive species policies could lead to inaccurate estimates of the invasion risk of

such taxa (Gordon et al. 2016). Economic losses arising from regulating taxa with high ornamental values can cause conflicts of interests among stakeholders, industries, and regulators (Wirth et al. 2004). As such, there is often a demand to develop and exempt infra- or interspecific taxa that are "safe" or "non-invasive" (Freyre et al. 2014; Guo et al. 2004).

There have been various attempts to develop sterile cultivars (e.g., Brand et al. 2012; Spies and du Plessis 1987; Wilson and Mecca 2003; reviewed by Datta et al. 2020) via methods such as genetic modification (Kanaya et al. 2008; Mitsuda et al. 2006), inducing polyploidy (Thammina et al. 2011), and interploid hybridization (Czarnecki et al. 2012; Deng et al. 2020). Czarnecki et al. (2012) and Deng et al. (2020) successfully bred and recommended various sterile cultivars of Lantana (*Lantana camara* L.) for ornamental use based on their relatively low seed production and viability. Other examples of sterile cultivars of invasive plants include Japanese spirea (*Spiraea japonica* L. f) (Wilson and Hoch 2009), Norway maple (*Acer platanoides* L.) (Conklin and Sellmer 2009), and heavenly bamboo (*Nandina domestica* Thunb.) (Knox and Wilson 2006). Even though these taxa were considered safe and non-invasive, this might change with plant age; for example, presumed sterile cultivars of Japanese barberry (*Berberis thunbergii* DC.) that were initially seedless started to produce seeds when plants were much older (Brand et al. 2012). Further, cultivars that have significantly lower seed germination and viability percentages need not necessarily have lower population growth rates (Knight et al. 2011; Le Roux et al. 2020; Wansell et al. 2022). Therefore, it is crucial that sterility is comprehensively assessed before deeming any infra- or interspecific entity as sterile.

Frameworks and protocols have been developed to identify safe/non-invasive cultivars of invasive plant species. Datta et al. (2020) framed a set of six questions that must be answered before a cultivar or hybrid is deemed safe. These six questions incorporate the main components of a risk analysis (risk identification, risk assessment, risk management, and risk communication) (Kumschick et al. 2020a, 2020b). Another example is the Intraspecific Taxon Protocol (ITP), a science-based assessment tool developed by researchers in Florida, USA, to assess cultivars with the potential of reduced invasiveness (IFAS 2008; Knox 2008). This tool contains a series of questions, the responses to which provide evidence that the cultivar: (1) can be easily distinguished from the wild-type species, (2) has traits that could reduce dispersal and/or spread, (3) is incapable of hybridizing with native flora, (4) does not readily revert to a natural or invasive form, and (5) is likely to have a lowered ecological impact. Applying the ITP protocol, the cultivars of *N. domestica* 'Fire Power' (Knox 2008; Knox and Wilson 2006) and 'Harbour Dwarf' (Knox and Wilson 2006) were deemed safe for Florida. Tools such as the Datta et al. (2020) framework and the ITP protocol can assist with developing regulations for infra and interspecific entities.

Very few regions have considered infra- or interspecific entities for invasive taxa regulations. In the United States, many states adopt independent procedures and protocols to identify and assess the impact of invasive species (Beaury et al. 2021; Lakoba et al. 2020). In Florida, as discussed previously, the ITP protocol is used. In Oregon, sterile cultivars can be approved for statewide sale if their seed production is less than 2%; however, a fee must be paid to Oregon State University to conduct a study to evaluate the fecundity of a specific taxon (Culley 2016), and the results must be submitted to the Oregon Department of Agriculture for verification. In Minnesota and Wisconsin, decisions on cultivar

**Table 1.** The definition of sterility in plants can vary with the context and goal for which the term is used, thus it is crucial that sterility is accurately assessed and defined when the term is being used.

Term	Definition
Sterile	The term “sterile,” when used within the context of plants in invasion science, relates to plants with poor quantity or quality of pollen and seeds (Czarnecki <i>et al.</i> 2012; Deng <i>et al.</i> 2020), which limits or prevents sexual reproduction. This definition does not preclude asexual reproduction, and so plants that are sterile under this definition can still form invasive populations.
Presumed sterile	Anecdotal evidence from the South African horticulture industry suggests that the term “sterile” was used to define a plant that is unable to escape from cultivation (captive/garden environment). Where this definition of “sterile” is used in the paper, we use the term “presumed sterile.”

bans or acceptability are based on scientific data pertaining to specific cultivars (Brand *et al.* 2012). Such approaches are backed up by scientific evidence, providing more confidence in the regulatory decisions, and thus results in increased research conducted to acquire evidence for safe/non-invasive cultivars within a specific region (Wilson and Deng 2023). Other states adopt different approaches, such as a decision-making tree to underpin cultivar exemption in New York (New York State Code of Rules and Regulations, Part 575, 2014) and the formation of a committee to explore sterile cultivar exemptions in Massachusetts, both of which are still evidence based (Brand 2016).

These examples show that the level of sterility deemed acceptable, and in fact how sterility is defined, varies. For example, anecdotal evidence from the South African horticulture industry suggests that the term “sterile” was used to define a plant that is unable to escape from cultivation, but still potentially able to produce viable seeds; therefore, we use the term “presumed sterile” here (see Table 1). The National Environmental Management: Biodiversity Act, Alien & Invasive Species (NEM:BA A&IS) regulations (Department of Environmental Affairs [DEA] 2014; Department of Forestry, Fisheries and the Environment [DFFE] 2020) attempt to accommodate infra- and interspecific variation in plant taxa by granting exemptions for cultivars or hybrids based on sterility. However, the evidence for exempting these “presumed sterile” cultivars and hybrids has not been published (Wilson and Kumschick 2024).

In this article, we first examine different approaches to regulate cultivars and hybrids of invasive plant species; second, we review how cultivars and hybrids of invasive plant species could be regulated by using South Africa as a case study; and finally, we provide guidelines for assessing sterility in cultivars/hybrids of invasive plant species.

### Approaches for Regulating Cultivars and Hybrids of Invasive Plant Species

For the regulation and subsequent management of invasive species, Kumschick *et al.* (2012) suggest that a transparent process is needed, which clearly expresses all the options by identifying and discussing the pros and cons of each. We present six approaches to regulate cultivars and hybrids of invasive plant species: (1) Globally Guilty by Association; (2) Nationally Guilty by Association; (3) Guilty until Proven Innocent; (4) Negotiated Guilt; (5) Claimed to Be Innocent; and (6) Innocent until Proven Guilty. The rationale, predicted number of entities banned, evidence required,

expected number of listing errors, and ease of implementation are shown for each approach in Table 2.

### The Rationale and Process Required for the Regulatory Approaches

The approaches range from risk adverse (Globally Guilty by Association) to reactive (Innocent until Proven Guilty). As such, they differ in the degree to which extrapolations of risk are made, resulting in a different risk assessment process for each approach. For the first two approaches “invasive elsewhere” is the only type of evidence required to ban cultivars or hybrids of an invasive taxon, thus no formal risk assessment process is required, rather these approaches are based on the precautionary principle with slightly different implications (see Table 2). While evidence based approaches 3 (Guilty until Proven Innocent) and 4 (Negotiated Guilt) still initially ban all cultivars and hybrids of the regulated invasive taxa; however, cultivars and hybrids of the regulated invasive taxa are allowed to be exempted from regulations if the required evidence is made available. The third approach requires scientific experiments to be conducted to demonstrate the safety of the cultivar or hybrid (this demonstration is usually in terms of sterility, see discussion on this later), which should feed into a formal risk analysis process for the exemption to be recommended. The fourth approach, also evidence based, does not require experiments to be conducted to demonstrate non-invasiveness, instead, the level of “guilt,” or the invasive potential of a cultivar, is negotiated between the regulators and stakeholders (i.e., a formal risk analysis might not be required) (Table 2). The fifth approach (Claimed to be Innocent) requires no evidence, and cultivars or hybrids can be exempted if any stakeholders claim that the entity is non-invasive (see Table 2 for the implications of such an approach). The last approach (Innocent until Proven Guilty) also does not require any evidence for safety. No formal risk assessment or risk analysis is required; all cultivars and hybrids of the regulated invasive taxa are exempted, unless there is specific evidence of invasiveness of the cultivar or hybrid. As such, this approach can be seen as reactive (Table 2).


### The Implications of the Various Regulatory Approaches

Each approach has different implications, as outlined in Table 2, and approaches differ in how easy they are to implement. The most resource-intensive approaches are those that require specific risk analyses to be conducted (approaches 3 and 4, Table 2). If cultivars or hybrids are granted exemptions, it is important that those entities can be easily distinguished from their parental genotypes in practice, thus avoiding confusion when such exemptions are implemented. It is also important to note that each approach would vary in other factors, such as number of entities banned, the effort required to demonstrate safety, and the expected number of listing errors (as outlined in Table 2), all of which are important aspects to consider when selecting a regulatory approach to implement.

Given invasiveness and impacts have some phylogenetic signal (Diez *et al.* 2012), we suggest that a reactive approach would be extremely risky, especially for cultivars or hybrids that have known invasive parent genotypes or congeners. On the other end of the spectrum, adopting the Globally Guilty by Association or Nationally Guilty by Association approach would likely mean a high number of infra- and interspecific taxa were unnecessarily banned, as risk assessments are not done at the infra- or interspecific taxonomic level (Gordon *et al.* 2016). Such approaches could lead to

**Table 2.** Six approaches for regulating cultivars and hybrids of invasive species from precautionary to reactive.

No.	Approach	Explanation of approach	Evidence required	Number of entities banned	Effort to demonstrate safety	Ease of implementation	Expected number of listing errors
1	Globally Guilty by Association	All cultivars or hybrids related to any species that are invasive anywhere in the world are regulated. It is based on the principle that the consequence of invasion is far higher than commercial interests. It is also consistent with the view that it is not possible to prove a particular genetic entity has, in perpetuity, lost the ability to become invasive.	Demonstrated invasiveness of the cultivar or hybrid relative anywhere in the world	High	Low	Fairly simple, providing taxa related to the entity can be identified	High number of taxa banned that are low risk (false positives)
2	Nationally Guilty by Association	If any entity is known to be invasive in the country adopting the regulation, all related cultivars and hybrids are banned. It is based on the principle that the consequence of invasion is far higher than commercial interests. It is also consistent with the view that it is not possible to prove a particular genetic entity has, in perpetuity, lost the ability to become invasive.	Demonstrated invasiveness of the cultivar or hybrid's relative in the same country adopting the regulation	High	Low	Fairly simple, providing taxa related to the entity can be identified	High number of taxa banned that are low risk (false positives)
3	Guilty until Proven Innocent	All entities are banned, but cultivars or hybrids can be exempted if evidence from experiments demonstrates an acceptable level of risk of a harmful invasion. It is the responsibility of the industry to demonstrate non-invasiveness.	Robust scientific experiments with large sample size and proper statistical analysis. Similar to prerelease safety assessment of biocontrol agents	High, but also dependent on number of tests done and the results of those tests	High	Difficult Time and resource intensive	Initially high number of taxa banned that do not need to be (false positives); however, allows for this number to be reduced over time
4	Negotiated Guilt	The level of guilt is set by agreements between regulators and users in the absence of strong information about invasiveness. Responsibility with regulator to demonstrate invasiveness OR industry to report invasiveness. An independent body, e.g., associated with the green industry, might be made responsible for approving entities based on known mechanisms and observations.	Primarily based on observations, potentially from multiple sources. Ideally, putative mechanism for sterility or non-invasiveness of cultivar/hybrid stated	Medium	Medium	Medium Negotiations can be lengthy and time-consuming; however, it is an inherently consultative and inclusive approach	Possibly low number of taxa banned that do not need to be (false positives) and taxa allowed that are high risk (false negatives)
5	Claimed to be Innocent	A cultivar or hybrid is exempted when any stakeholder claims it to be non-invasive. It is based purely on anecdotal observations and good faith	No evidence required	Low	Low	Few resources needed	Possibly high number of taxa allowed that are high risk (false negatives)
6	Innocent until Proven Guilty	All cultivars or hybrids are exempt from regulations unless there is evidence for invasiveness. This approach argues that cultivars or hybrids differ significantly from the invasive species and therefore the invasiveness of the cultivar or hybrid must first be proved/demonstrated before it can be banned.	Responsibility with regulator to provide scientific evidence demonstrating invasiveness	Low	Low	No resources required for cultivars or hybrids to be regulated. However, evidence required to prove invasiveness; costs of producing such evidence possibly lies with society	Highest number of possible taxa allowed which are high risk (false negatives)



**Figure 1.** Photo panel illustrating examples of “presumed sterile” cultivars in South: (A) *Duranta erecta* ‘Sapphire Showers’; (B) *Duranta erecta* ‘Sheena’s Gold’; (C) *Duranta erecta* ‘Goldmine’; (D) *Vinca major* ‘Variegata.’ (Photos by Duran Chetty).

significant economic losses in the green industry, potentially generating substantial disputes between the green industry and regulators, particularly if there was no clear route to contest the listing of taxa that stakeholders perceive to be safe.

### Regulation of “Presumed Sterile” Cultivars and Hybrids in South Africa

In the following section, we used South Africa as a case study to review the regulation of “presumed sterile” cultivars and hybrids of invasive taxa. There have been various negotiations between the South African horticultural industry and the South African Department of Forestry, Fisheries and the Environment (DFFE) regarding the A&IS Regulations and “presumed sterile” cultivars and hybrids (K Montgomery, personal communication).

Because all of the “presumed sterile” (discussed later) cultivars or hybrids registered in South Africa (see Supplementary Table S1) are important ornamental plants (see Figure 1 for examples) or horticultural trees (Armitage 2008; Gardening in South Africa 2023), the horticulture industry is a crucially interested and affected stakeholder. Thus, the South African horticulture industry initiated negotiations with the government in 2001 to prevent the regulation of taxa that were horticulturally and economically significant (Figure 2). The negotiations entailed discussions regarding the invasiveness (or lack thereof) of those taxa before implementation of the regulations.

In 2004, the National Environmental Management: Biodiversity Act (NEM:BA) (Act No. 10 of 2004) was enacted (see Lukey and Hall [2020] and Wilson and Kumschick [2024] for a detailed review of the history of invasive species regulations in South Africa). As part of the NEM:BA, Alien and Invasive Species (A&IS) Regulations were promulgated in 2014 ([DEA

Department of Environmental Affairs, 2014). Under revised lists of October 2020 (DFFE 2020), 382 plant taxa were listed (Wilson 2024). In 2004, when NEM:BA was enacted, the South African horticulture industry requested exemptions for cultivars and hybrids for specific taxa; as such, the NEM:BA A&IS Regulations address infra- and interspecific variation by granting exemptions for cultivars or hybrids based on their presumed sterility.


On this basis, the negotiations moved toward exemptions of cultivars and hybrids that were seemingly safe. These negotiations were lengthy, and it was decided in 2010 (Figure 2) to implement the “polluters pay” principle, suggesting that those responsible for causing harm to the environment should be responsible for the cost of such damage (Luppi *et al.* 2012), increasing conflict between the negotiating parties. It was agreed that a consensus had to be reached, but there was still a lack of scientific evidence regarding the invasiveness of the debated taxa.

Eventually, a consensus was reached in 2014. As part of this consensus, and before the first A&IS Lists were promulgated in 2014 ([DEA] Department of Environmental Affairs, 2014), the DFFE gave the industry an opportunity to propose a list of those cultivars and hybrids that it regarded to be sterile (Figure 2, “the unofficial consensus list”; see Supplementary Table S1). This list contained 187 cultivars and/or hybrids from 34 taxa (see Supplementary Table S1). This “presumed sterile” taxa list has changed over time (Figure 2) and currently contains 157 “presumed sterile” cultivars and/or hybrids from 16 taxa (Table 3; Supplementary Table S2). However, the evidence for the sterility of the registered cultivars and hybrids is lacking. Rather, the argument for the requested exemptions was on the basis that none of the cultivars nor hybrids had recorded naturalized populations. For legislative purposes, the government termed these “presumed sterile” taxa as sterile cultivars and hybrids, bringing forth the inception of the term “sterile cultivars and hybrids” in the South African context (see Table 1). However, the industry and DFFE did not publish evidence of sterility for “presumed sterile” cultivars and hybrids (see Table 4 for anecdotes from the green industry regarding “presumed sterile” cultivars).

In 2017, revisions to the unofficial consultative list included the removal of tickseed (*Coreopsis lanceolata* L.) cultivars due to a lack of evidence of sterility, and the removal of greater periwinkle (*Vinca major* L.) cultivars due to the plant spreading more vegetatively than sexually (Figure 2; Supplementary Appendix). The industry was given deadlines (1 to 2 years) to prove the sterility of various listed cultivars and hybrids if they were to remain on the unofficial consultative list (Supplementary Appendix). In the 2020 iteration of the NEM:BA A&IS Lists (DFFE 2020), less than half (16 out of 34) taxa still had regulatory provisions for exemptions of “presumed sterile” cultivars (Figure 2; Table 3; Supplementary Table S2). However, we are not aware of publicly available evidence of sterility for the retained entities or of the precise reason why specific entities were removed from the lists. Thus, it is important to understand how the list proposed by the South African horticulture industry was developed so a transparent, evidence-based approach can be used in future for producing lists that guide policy and action (Butchart *et al.* 2010; Perry and Perry 2008).

### What Should a Sterility Assessment for Cultivars or Hybrids of Invasive Species Include?

Here, we outline the basic requirements for sterility assessments of cultivars/hybrids of invasive plant species and present a generic pro forma for reporting on sterility (Figure 3). It is important to note



**Figure 2.** Timeline displaying the major events of the negotiations between the South African Horticultural Industry (SAHI) and the Department of Forestry, Fisheries and the Environment (DFFE) regarding “presumed sterile” cultivars of invasive plants in South Africa. For a detailed timeline of the NEM:BA A&IS Regulations and Lists, see Wilson and Kumschick (2024).

that the guidelines presented are not a standardized protocol for sterility assessments but aim to outline various components that should be included in any sterility assessments for invasive plants and to provide examples of the types of experiments that could be conducted to gather the required data. The first three components of the sterility assessment specifically deal with assessing the sexual reproductive pathway, the fourth with quantifying asexual reproduction, and the last component assesses the stability of sterility.

### Flower and Fruit Production

Do the cultivars/hybrids produce flowers and fruit, and if so, how many? Common-garden or greenhouse experiments (e.g., Knox and Wilson 2006) can be set up by growing replicates of each tested cultivar/hybrid for a period of time (until reproductive maturity). Wild-type plants can be grown as controls.

### Pollen Analyses

If the tested cultivars/hybrids produce flowers, pollen viability analyses should be done. Pollen viability assessments are often done using biological staining techniques (Jones 2012; Pinillos and Cuevas 2008) or by conducting pollen germination experiments. For the control of these experiments, pollen from wild-type plants (representing the invasive forms) of the tested cultivar/hybrids should be used, and the pollen viability and/or germination percentages should be significantly lower in tested hybrids/cultivars than in the wild type. If no flowers are produced, then the cultivar/hybrid can be defined as truly sterile. However, if pollen viability/germination is below a certain percentage threshold this can mean a taxon is not invasive. It should be up to the legislators and stakeholders within a region/country to agree on an

acceptable threshold, as this can vary between various taxa depending on other factors such as time to reproduction (different in fast- and slow-growing species), benefits of cultivar (environmental, economic, or social) and results from seed analyses (step 3). Finally, electron microscopy can be used to supplement pollen viability results, which may identify abnormal pollen grains characteristic of low pollen viability (Shaik et al. 2023).

### Seed Analyses

If cultivars/hybrids produce fruit (and subsequently set seed), the number of seeds per fruit should be quantified. Thereafter, seed viability and germination assays (e.g., Czarnecki et al. 2012; Deng et al. 2020) should be conducted. Seed viability assays can be done using the standard tetrazolium test, and germination assays can be conducted *in vitro* (petri dishes and incubators) or *ex vitro* (sowing in soil). A cultivar/hybrid that does not produce seeds or whose seeds have 0% viability or germination can be defined as truly sterile. However, as per the previous step, an acceptable threshold can be defined, which may be context specific. Various other/additional seed analyses can be done (seed mass/ultra-structure analyses, etc.) to understand the mechanisms of sterility.

These three components provide insights into the degree of fertility of a cultivar or hybrid. The next two steps aim to provide a more comprehensive understanding of the invasion risk.

### Potential for Vegetative Propagation

Can the cultivars/hybrids reproduce asexually? Experiments can be conducted to determine the survival/success rate of propagation from cuttings (or other types of vegetative propagation, such as bulbils) to gain an understanding of

**Table 3.** Taxa listed under the 2014 and 2020 National Environmental Management: Biodiversity Act, Alien and Invasive Species Regulations for which there is or was provision to exempt sterile cultivars or hybrids<sup>a</sup>.

scientificName <sup>b</sup>	Other names used <sup>c</sup>	vernacularName <sup>d</sup>	Number of “presumed sterile” entities exempted	
			2014	2020
<i>Acer negundo</i> L.	None	Ash-leaved maple, box elder	0 <sup>e</sup>	0 <sup>e</sup>
<i>Ageratum houstonianum</i> Mill.	None	Mexican ageratum	6	6
<i>Berberis thunbergii</i> DC.	None	Japanese barberry	10	10
<i>Buddleja davidii</i> Franch.	None	Chinese sagewood, summer lilac	7	7
<i>Canna indica</i> L.	None	Indian shot	0	No provisions
<i>Catharanthus roseus</i> (L.) G. Don	None	Madagascar periwinkle	77	77
<i>Cenchrus setaceus</i> (Forssk.) Morrone	<i>Pennisetum setaceum</i> (Forssk.) Chiov.	Fountain grass	5	5
<i>Cestrum</i> L.	None	<i>Cestrum</i> spp.	0	No provisions
<i>Coreopsis lanceolata</i> L.	None	Tickseed	0	No provisions
<i>Cortaderia selloana</i> (Schult. & Graebn.) Asch. & Graebn.	None	Pampas grass	2	No provisions
<i>Duranta erecta</i> L.	<i>Duranta repens</i> L., <i>Duranta plumieri</i> Jacq.	Forget-me-not-tree, pigeon berry	6	6
<i>Gleditsia triacanthos</i> L.	none	Honey locust	6	No provisions
<i>Hedera canariensis</i> Willd.	<i>Hedera helix</i> L. subsp. <i>canariensis</i> (Willd.) Cout.	Canary ivy, Madeira ivy, Algerian ivy	3	3
<i>Hedera helix</i> L.	<i>Hedera helix</i> L. subsp. <i>helix</i>	English ivy	20	20
<i>Ipomoea indica</i> (Burm.) Merr.	<i>Ipomoea congesta</i> R.Br.	Blue morningglory	0	No provisions
<i>Ipomoea purpurea</i> (L.) Roth	None	Purple morningglory	0	No provisions
<i>Ligustrum lucidum</i> W.T. Aiton	None	Chinese wax-leaved privet	1	No provisions
<i>Ligustrum ovalifolium</i> Hassk.	None	Californian privet	1	1
<i>Limonium sinuatum</i> (L.) Mill.	None	Statice, sea lavender	11	No provisions
<i>Metrosideros excelsa</i> Sol. ex Gaertn.	<i>Metrosideros tomentosa</i> A. Rich.	New Zealand Christmas tree	6	No provisions
<i>Melaleuca viminalis</i> (Sol. ex Gaertn.) Byrnes	<i>Callistemon viminalis</i> (Sol. ex Gaertn.) G. Don	Weeping bottlebrush	6	6
<i>Morus alba</i> L.	None	White mulberry, common mulberry	0	No provisions
<i>Murraya paniculata</i> (L.) Jack	<i>Murraya exotica</i> L.	Orange jessamine	1	1
<i>Nephrolepis cordifolia</i> (L.) C. Presl	<i>Polypodium cordifolium</i> L.	Erect sword fern, ladder sword fern	2	No provisions
<i>Nephrolepis exaltata</i> (L.) Schott	<i>Polypodium exaltatum</i> L.	Sword fern, Boston sword fern	11	11
<i>Nerium oleander</i> L.	None	Oleander	1 + all double flowering cultivars	No provisions
<i>Opuntia ficus-indica</i> (L.) Mill.	<i>Opuntia megacantha</i> Salm-Dyck	Mission prickly pear, sweet prickly pear	4	4
<i>Opuntia robusta</i> H.L. Wendl. ex Pfeiff.	None	Blue-leaf cactus	0	0
<i>Pyracantha angustifolia</i> (Franch.) C.K. Schneid.	None	Yellow firethorn	0	No provisions
<i>Pyracantha coccinea</i> M. Roem.	None	Red firethorn	0	No provisions
<i>Pyracantha crenulata</i> (D. Don) M. Roem. <sup>f</sup>	<i>Pyracantha crenatoserrata</i> (Hance) Rehder, <i>Pyracantha fortuneana</i> (Maxim.) H.L.Li, <i>Pyracantha crenulata</i> (D. Don) M. Roem. var. <i>rogersiana</i> , <i>Pyracantha rogersiana</i> (A.B. Jacks.) Chitt.	Chinese firethorn, broad leaf firethorn, Himalayan firethorn	0	No provisions
<i>Pyracantha koidzumii</i> (Hayata) Rehder	None	Formosa firethorn	0	No provisions
<i>Vinca major</i> L.	None	Greater periwinkle	2	No provisions
<i>Vinca minor</i> L.	None	Lesser periwinkle	4	4

<sup>a</sup>Column headings are as per DarwinCore terms where available. For full details, see Supplementary Tables S1 and S2.

<sup>b</sup>The scientificName was taken from Wilson (2024), with nomenclature checked therein against the Botanical Database of Southern Africa (BODATSA) and Plants of the World Online (POWO) during 2023.

<sup>c</sup>Other names used are either synonyms used in at least one version of the regulatory lists, synonyms specified in the regulatory listing [e.g., the NEM:BA A&S List includes the following listing: “*Duranta erecta* L. (= *D. repens* L., *D. plumieri* Jacq.)”], or names misapplied in South Africa specified in the regulations (e.g., *Pyracantha fortuneana* was misapplied to *Pyracantha crenulata*).

<sup>d</sup>The vernacularName is as presented exactly in the NEM:BA A&S Lists (including capitalization).

<sup>e</sup>It is unclear if there were any submissions of sterile cultivars/hybrids for *Acer negundo*, as this taxon does not appear in the unofficial consultative list (Supplementary Table S1); however, there have been provisions for exemptions for this taxon since 2014.

<sup>f</sup>*Pyracantha crenatoserrata* is a recognized synonym of *Pyracantha crenulata* (Plants of the World Online) but is listed separately in the NEM:BA A&S Lists.

**Table 4.** Anecdotes from the green industry regarding “presumed sterile” cultivars or hybrids<sup>a</sup>.

Plant/term	Discussion
Variegated forms	A prevalent notion is that variegated forms of known invasive species are sterile. For example, the variegated form of <i>Vinca major</i> was considered sterile and suitable for trade (2014–2017) in South Africa; however, this was amended, as the cultivar was observed to spread vegetatively. Variegated forms might fix less carbon than nonvariegated forms and so have slower growth rates, but we know of no evidence that confirms sterility in variegated forms.
Bonsai plants	Due to small stature, the fecundity of the bonsai plants is limited. Bonsai plants are usually maintained indoors or in a highly managed environment, unlike many garden plants that can occur on the edges of gardens or are not well maintained. As such, the possibility of a bonsai becoming invasive is very low. However, in principle, even a few viable seeds could lead to the start of a new population. Because dwarfness in bonsai is usually human induced rather than genetic, the progeny from bonsai can become full-sized trees.
Purple fountain grass ( <i>Pennisetum × advena</i> ‘Rubrum’)	Fountain grass is regulated in South Africa, but the red/purple cultivars have very high ornamental value and are considered sterile. They are still sold in the nurseries as a “presumed sterile” cultivar. However, we know of no published evidence demonstrating the sterility of this cultivar. The issue of “presumed sterile” cultivars pertains not only to cultivars of <i>Pennisetum × advena</i> in South Africa but extends to 15 other taxa (as shown in Table 1), encompassing a total of 157 “presumed sterile” cultivars/hybrids in South Africa.

<sup>a</sup>It is interesting to note existing impressions that among members of the green industries regarding certain cultivars and hybrids perceived to be “non-invasive” and safe for trade. These perceptions are anecdotal evidence based on long-term observations. Some of these are briefly discussed in this table.

how easily the plants can propagate (or spread) asexually. Further, it is advised that basic vigor assessments be conducted with the surviving cuttings to determine which are the fastest-growing cultivars/hybrids (a trait that increases invasive potential).

### Potential for Genetic Changes

Finally, to determine the potential for genetic changes that could affect the stability of sterility, cross-breeding between cultivars and between cultivars and wild-type plants should be investigated. Hand-pollination experiments in the greenhouse are recommended (e.g., Wilson and Hoch 2009), and the seeds of the F<sub>1</sub> progeny should be tested as per component 3 of the sterility assessment. If seed production is significantly higher, then the risk assessment needs to be adjusted.

### Discussion

For the regulation of cultivars and hybrids of invasive taxa, South Africa initially adopted an approach similar to Claimed to be Innocent (Table 2, approach 5), but subsequently moved to something between the Negotiated Guilt (Table 2, approach 4), and Guilty until Proven Innocent (Table 2, approach 3). Adopting a Claimed to be Innocent approach is risky, and the approach may change over time, as seen in the case of South Africa, where taxa that were exempt, such as *V. major*, among others, were later banned (see Supplementary Appendix for other examples). Hence, we primarily recommend a Guilty until Proven Innocent approach for regulating cultivars or hybrids of invasive taxa at a national scale (i.e., all cultivars or hybrids, where at least one related taxon is regulated should be regulated unless there is documented evidence that the cultivar/hybrid is not invasive). Sterile cultivars or hybrids of invasive plant species could be ideal candidates for such exemptions, but it is crucial that within the supporting evidence, sterility is not only appropriately assessed but also accurately defined.

Allowing for the independent regulation of cultivars and hybrids of invasive plant species by exempting safe/non-invasive (usually synonymous with sterile) taxa that are underpinned by scientific evidence may be the most viable regulatory option. However, there will still remain other risks, such as the

misidentification of sterile cultivars or hybrids. Traders may knowingly or unknowingly label specific cultivars or hybrids with the names of the exempted taxa. Thus, it is recommended that procedures are put in place to help prevent this, such as routine genetic testing (e.g., DNA fingerprinting or sequencing approaches) and plant auditing. Further, public (and nursery customer) awareness would be needed for the approach to be widely acceptable and adopted. If the recommended regulatory approach is adopted nationally, there would be an incentive to patent sterile plants, making plant tracking easier. Finally, generic pro forma for reporting sterility based on observations and/or experiments (such as that presented in Figure 3) may assist with making the regulatory approach easier to implement.

### Recommendations

For the case of regulating cultivars or hybrids, we primarily recommend evidence-based approaches such as the Guilty until Proven Innocent or the Negotiated Guilt approaches. The Guilty until Proven Innocent approach is the most time-consuming and stringent, resulting in a relatively high number of taxa being banned, but it is also the most evidence-based approach that seeks to minimize conflict of interest between stakeholders and legislators. This approach can be justified for regulating cultivars or hybrids, because such taxa generally display similar traits to their parent genotypes; thus, the precautionary principle still applies. Although this approach requires substantial information, it still allows for exemptions of cultivars or hybrids based on evidence gained from robust scientific experiments, making provision for “safe” taxa claims. This type of approach has been demonstrated in Oregon, USA, where 18 sterile cultivars of the invasive Chinese sagewood (*Buddleja davidii* Franch.) were deemed safe for trade/use (underpinned by scientific evidence), which served as a model for other states to follow (Contreras and McAninch 2013). Further, adopting a Guilty until Proven Innocent approach encourages research, which increases the output and volume of scientific knowledge pertaining to safe/non-invasive cultivars, as seen in Florida (Wilson and Deng 2023). In the event that an Innocent until Proven Guilty approach cannot be successfully adopted, we recommend a Negotiated Guilt approach. This approach does not require scientific evidence or assessments,

Pro- forma (template): **What should a sterility assessment for cultivars or hybrids of invasive species include?**

**1a. Does the entity produce flowers?** (Yes/No/Not studied/NA)

Notes on evidence: examples include long-term flower evaluation with greenhouse or common-garden experiments.

**1b. Does the entity produce fruit?** (Yes/No/Not studied/NA)

Notes on evidence: examples include long-term fruit evaluation with greenhouse or common-garden experiments

**2. Does the entity produce viable pollen?** (Yes/No/Not studied/NA - if no flowers)

If yes, how does production and/or viability compare with other entities?

Notes on evidence: examples include pollen viability and germination experiments.

**3. Does the entity produce viable seeds?** (Yes/No/Not studied/NA)

If yes, how does production and/or viability compare with other entities?

Notes on evidence: examples include seed viability and germination experiments.

**4. Does the entity reproduce vegetatively?** (Yes/No/Not studied/NA)

If yes, how does it compare to other entities?

Notes on evidence: examples include cuttings propagation experiments along with vigour and growth rate assessments.

**5. Does the entity become fertile/remain sterile after out-crossing?** (Yes/No/Not studied/NA):

Notes on evidence: examples include cross-pollination experiments to determine seed set and seed viability.

**Overall conclusion**

Example: The entity is dependent on seed production for it to become invasive. The entity produces fertile seeds but at a much lower level than parental stock. The likelihood and rate of an invasion of the entity is therefore likely to be substantially lower than parental stock but it is not clear that the overall risk is below an acceptable threshold.

**Figure 3.** A generic pro forma for reporting sterility based on observations and/or experiments on: flowering, fruiting, pollen, and seeds; the potential for vegetative propagation; and the potential for genetic changes (including hybridization and reversion to fertility).

exemptions are based on known mechanisms and long-term observations that an independent body associated with the green industry is responsible for. However, it must be noted that the Negotiated Guilt approach may lead to lengthy negotiations between regulators and users or inaccurate risk assessments of the exempted taxa.

## Conclusion

Regulating infra- and interspecific entities of taxa that are known to be invasive can be a complex task but is important if conflicts of interest between various stakeholders (primarily from industry) and regulators are to be resolved. South African regulation of infra- and interspecific entities of invasive taxa is laudable, but the process can be more transparent and evidence based. We recommend that for any country with cultivars or hybrids of non-native plant species, exemptions for infra- and interspecific entities should be on the basis of risk analyses for those entities (e.g., Kumschick et al. [2020b] for South Africa). Ideally risk analyses of the related entities should also be produced with clear explanations as to why the risk differs. We believe such exemptions should be specified in official documentation, so they are transparent.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/inp.2024.24>

**Acknowledgments.** South African Department of Forestry, Fisheries and the Environment (DFFE) officials are thanked for their input regarding the regulations. Kay Montgomery is thanked for the valuable information provided pertaining to the South African horticultural industry. Katelyn Faulkner, Ashlyn Padayachee, Michael Cheek and two anonymous reviewers are thanked for comments on the article.

**Funding statement.** The South African Department of Forestry, Fisheries and the Environment (DFFE) is thanked for funding, noting that this publication does not necessarily represent the views or opinions of DFFE or its employees. DC acknowledges the Cape Peninsula University of Technology for research support. SK acknowledges the support of the Centre for Invasion Biology (CIB) at Stellenbosch University.

**Competing interests.** The authors declare no competing interests.

## References

- [DEA] Department of Environmental Affairs (2014) National Environmental Management: Biodiversity Act (10/2004): Alien and Invasive Species Regulations, 2014. Government Gazette, notice 598 of 2014, no. 37885, August 1, 2014, Pretoria, South Africa. Pp 3–32
- [DFFE] Department of Forestry, Fisheries and the Environment (2020) National Environmental Management: Biodiversity Act (Act No. 10 of 2004): Alien and Invasive Species Regulations. Government Gazette, notice 1020 of 2020, no. 43735, September 25, 2020, Pretoria, South Africa. Pp 14–44
- [IPBES] Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (2023) Summary for Policymakers of the Thematic Assessment Report on Invasive Alien Species and Their Control of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Bonn, Germany: IPBES Secretariat. 952 p
- Armitage A (2008) A Great Group of Plants: Pigeon Berries, a.k.a. *Duranta erecta*. Greenhouse Grower. <http://www.greenhousegrower.com/crops/a-great-group-of-plants-pigeon-berries-a-k-a-duranta-erecta>. Accessed: June 10, 2023
- Beaury EM, Fusco EJ, Allen JM, Bradley BA (2021) Plant regulatory lists in the United States are reactive and inconsistent. *J Appl Ecol* 58:1957–1966
- Brand M (2016) Sterile cultivars (or close to it)—is this a viable option for the nursery industry? *Acta Hort* 1140:273–278
- Brand MH, Lehrer JM, Lubell JD (2012) Fecundity of Japanese barberry (*Berberis thunbergii*) cultivars and their ability to invade a deciduous woodland. *Invasive Plant Sci Manag* 5:464–476
- Buhk C, Thielsch A (2015) Hybridisation boosts the invasion of an alien species complex: Insights into future invasiveness. *Perspect Plant Ecol Evol Syst* 17:274–283
- Butchart SH, Walpole M, Collen B, van Strien A, Scharlemann JP, Almond RE, Baillie JE, Bomhard B, Brown C, Bruno J, Carpenter KE, Carr GM, Chanson J, Chenery AM, Csirke J, et al. (2010) Global biodiversity: indicators of recent declines. *Science* 32:1164–1168
- Charlesworth B (2009) Effective population size and patterns of molecular evolution and variation. *Nat Rev Genet* 10:195–205
- Conklin JR, Sellmer JC (2009) Flowering, fecundity, seed germination, and seed viability of *Viburnum opulus* L. cultivars. *J Environ Hort* 27:31–36
- Contreras R, McAninch G (2013) Back from the ban: new *Buddleja* cultivars prove it can make sense to legalize sterile versions of some invasives. *Digger* 57:33–36
- Culley TM (2016) Identifying invasive plant species: what plant propagators need to know about the science behind invasive plant assessment protocols. *Acta Hort* 1140:265–272
- Culley TM, Hardiman NA (2009) The role of intraspecific hybridization in the evolution of invasiveness: a case study of the ornamental pear tree *Pyrus calleryana*. *Biol Invasions* 11:1107–1119
- Czarnecki DM, Wilson SB, Knox GW, Freyre R, Deng Z (2012) UF-T3 AND UF-T4: two sterile *Lantana camara* cultivars. *HortScience* 47:132–137
- Datta A, Kumschick S, Geerts S, Wilson JR (2020) Identifying safe cultivars of invasive plants: six questions for risk assessment, management, and communication. *NeoBiota* 62:81–97
- Deng Z, Wilson SB, Ying X, Chen C, Freyre R, Zayas V, Czarnecki DM (2020) “UF-1013-1”: an infertile cultivar of *Lantana camara*. *HortScience* 55:953–958
- Diez JM, Hulme PE, Duncan RP (2012) Using prior information to build probabilistic invasive species risk assessments. *Biol Invasions* 14:681–691
- Ellstrand NC, Schierenbeck KA (2000) Hybridization as a stimulus for the evolution of invasiveness in plants? *Euphytica* 148:35–46
- Freyre R, Moseley A, Knox GW, Wilson SB (2012) Fruitless *Ruellia simplex* R10-102 (“Mayan Purple”) and R10-108 (“Mayan White”). *HortScience* 47:1808–1814
- Freyre R, Wilson SB, Knox GW (2014) Breeding *Ruellia* and trialing for sterility at the University of Florida. *Acta Hort* 1055:431–435
- García-de-Lomas J, Vilà M (2015) Lists of harmful alien organisms: are the national regulations adapted to the global world? *Biol Invasions* 17:3081–3091
- Gardening in South Africa (2023) Home page. <http://www.gardeninginsouthafrica.co.za>. Accessed: May 30, 2023
- Gordon DR, Flory SL, Lieurance D, Hulme PE, Buddenhagen C, Caton B, Champion PD, Culley TM, Daehler C, Essl F, Hill JE, Keller RP, Kohl L, Koop AL, Kumschick S, et al. (2016) Weed risk assessments are an effective component of invasion risk management. *Invasive Plant Sci Manag* 9:81–83
- Gotelli NJ, Stanton-Geddes J (2015) Climate change, genetic markers and species distribution modelling. *J Biogeogr* 42:577–1585
- Guo WW, Prasad D, Cheng YJ, Serrano P, Deng XX, Grosse JW (2004) Targeted cybridization in citrus: transfer of Satsuma cytoplasm to seedy cultivars for potential seedlessness. *Plant Cell Rep* 22:752–758
- Hereford J (2009) A quantitative survey of local adaptation and fitness trade-offs. *Am Nat* 173:579–588
- Hovick SM, Whitney KD (2014) Hybridisation is associated with increased fecundity and size in invasive taxa: meta-analytic support for the hybridisation-invasion hypothesis. *Ecol Lett* 17:1464–1477
- Hulme PE, Brundu G, Carboni M, Dehnen-Schmutz K, Dullinger S, Early R, Essl F, González-Moreno P, Groom QJ, Kueffer C, Kühn I, Maurel N, Novoa A, Pergl J, Pyšek P, et al. (2018) Integrating invasive species policies across ornamental horticulture supply chains to prevent plant invasions. *J Appl Ecol* 55:92–98
- IFAS (2008) Invasive Plant Working Group. IFAS Intraspecific Taxon Protocol. Gainesville: University of Florida/IFAS, Center of Aquatic and Invasive Plants. <https://edis.ifas.ufl.edu/publication/AG376>. Accessed: May 25, 2023
- Jones GD (2012) Pollen analyses for pollination research, unacetolyzed pollen. *J Pollinat Ecol* 9:96–107

- Kanaya T, Saito H, Hayashi Y, Fukunishi N, Ryuto H, Miyazaki K, Kusumi T, Abe T, Suzuki KI (2008) Heavy-ion beam-induced sterile mutants of verbena (*Verbenax hybrida*) with an improved flowering habit. *Plant Biotechnol J* 25:91–96
- Knight TM, Havens K, Vitt P (2011) Will the use of less fecund cultivars reduce the invasiveness of perennial plants? *BioScience* 61:816–822
- Knox GW (2008) Invasive issues: how to determine which plants are invasive and where do cultivars fit? Pages 533–536 in *Combined Proceedings, Volume 58, International Plant Propagators' Society, Seattle, Washington*
- Knox GW, Wilson SB (2006) Evaluating north and south Florida landscape performance and fruiting of ten cultivars and a wildtype selection of *Nandina domestica*, a potentially invasive shrub. *J Environ Hortic* 24:137–142
- Kumschick S, Bacher S, Dawson W, Heikkilä J, Sendek A, Pluess T, Robinson TB, Kuhn I (2012) A conceptual framework for prioritization of invasive alien species for management according to their impact. *NeoBiota* 15:69–100
- Kumschick S, Fernandez Winzer L, McCulloch-Jones EJ, Chetty D, Fried J, Govender T, Potgieter LJ, Rapetsoa MC, Richardson DM, van Velden J, Van der Colff D, Miz S, Wilson JR (2024) Considerations for developing and implementing a safe list for alien taxa. *BioScience* 74:97–108
- Kumschick S, Foxcroft LC, Wilson JR (2020a) Analysing the risks posed by biological invasions to South Africa. Pages 573–595 in van Wilgen B, Measey J, Richardson D, Wilson J, Zengeya T, eds. *Biological Invasions in South Africa. Invading Nature—Springer Series in Invasion Ecology 14*. Cham, Switzerland: Springer
- Kumschick S, Richardson DM (2013) Species-based risk assessments for biological invasions: advances and challenges. *Divers Distrib* 19:1095–1105
- Kumschick S, Wilson JR, Foxcroft LC (2020b) A framework to support alien species regulation: the Risk Analysis for Alien Taxa (RAAT). *Neobiota* 62:213–239
- Lakoba T, Brooks RK, Haak DC, Barney JN (2020) An analysis of US state regulated weed lists: a discordance between biology and policy. *Bioscience* 70:804–813
- Landry CR, Hartl DL, Ranz JM (2007) Genome clashes in hybrids: insights from gene expression. *Heredity* 99:483–493
- Le Roux JJ, Clusella-Trullas S, Mokotjomela TM, Mairal M, Richardson DM, Skein L, Wilson JR, Weyl OL, Geerts S (2020) Biotic interactions as mediators of biological invasions: insights from South Africa. Pages 387–427 in van Wilgen B, Measey J, Richardson D, Wilson J, Zengeya T, eds. *Biological Invasions in South Africa. Invading Nature—Springer Series in Invasion Ecology 14*. Cham, Switzerland: Springer
- Leimu R, Fischer M (2008) A meta-analysis of local adaptation in plants. *PLoS ONE* 3:e4010
- Linhart YB, Grant MC (1996) Evolutionary significance of local genetic differentiation in plants. *Annu Rev Ecol Evol* 27:237–277
- Lukey P, Hall J (2020) Biological invasion policy and legislation development and implementation in South Africa. Pages 515–551 in van Wilgen BW, Measey J, Richardson DM, Wilson JR, Zengeya TA, eds. *Biological Invasions in South Africa. Invading Nature—Springer Series in Invasion Ecology 14*. Cham, Switzerland: Springer
- Luppi B, Parisi F, Rajagopalan S (2012) The rise and fall of the polluter-pays principle in developing countries. *Int. Rev Law Econ* 32:135–144
- McGeoch MA, Spear D, Kleynhans EJ, Marais E (2012) Uncertainty in invasive alien species listing. *Ecol Appl* 22:959–971
- Mitsuda N, Hiratsu K, Todaka D, Nakashima K, Yamaguchi-Shinozaki K, Ohme-Takagi M (2006) Efficient production of male and female sterile plants by expression of a chimeric repressor in Arabidopsis and rice. *Plant Biotechnol J* 4:325–332
- Mizutani T, Goka K (2010) Japan's Invasive Alien Species Act. *Appl Entomol Zool* 45:65–69
- Novoa A, Flepu V, Boatwright J (2018) Is spinelessness a stable character in cactus pear cultivars? Implications for invasiveness. *J Arid Environ* 160:11–16
- Pergl J, Sádlo J, Petrušek A, Laštůvka Z, Musil J, Perglová I, Šanda R, Šefrová H, Šíma J, Vohralík V, Pyšek P (2016) Black, Grey and Watch Lists of alien species in the Czech Republic based on environmental impacts and management strategy. *NeoBiota* 28:1–37
- Perry D, Perry G (2008) Improving interactions between animal rights groups and conservation biologists. *Conserv Biol* 22:27–35
- Pinillos V, Cuevas J (2008) Standardization of the fluorochromatic reaction test to assess pollen viability. *Biotech Histochem* 83:15–21
- Pyšek P, Jarošík V, Hulme PE, Pergl J, Hejda M, Schaffner U, Vilà M (2012) A global assessment of invasive plant impacts on resident species, communities and ecosystems: the interaction of impact measures, invading species' traits and environment. *Global Chang Biol* 18:1725–1737
- Shaik S, Chetty D, Watt MP (2023) Micropropagation decreases pollen and seed viabilities of two *Solanum nigrum* clonal genotypes. *S Afr J Bot* 153: 203–208
- Spies JJ, du Plessis H (1987) Sterile *Lantana camara*: fact or theory. *S Afr J Plant Soil* 4:171–174
- Thammina C, He M, Lu L, Cao K, Yu H, Chen Y, Tian L, Chen J, McAvooy R, Ellis D, Zhao D, Wang Y, Zhang X, Li Y (2011) In vitro regeneration of triploid plants of *Euonymus alatus* 'Compactus' (Burning Bush) from endosperm tissues. *HortScience* 46:1141–1147
- Thompson GD, Robertson MP, Webber BL, Richardson DM, Le Roux JJ, Wilson JR (2011) Predicting the subspecific identity of invasive species using distribution models: *Acacia saligna* as an example. *Divers Distrib* 17:1001–1014
- Tokarska-Guzik B, Bzdęga K, Dajdok Z, Mazurska K, Solarz W (2021) Invasive alien plants in Poland—the state of research and the use of the results in practice. *Environ Socio-Econ Stud* 9:71–95
- Turbelin AJ, Malamud BD, Francis RA (2017) Mapping the global state of invasive alien species: patterns of invasion and policy responses. *Global Ecol Biogeogr* 26:78–92
- Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V, Maron JL, Pergl J, Schaffner U, Sun Y, Pyšek P (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol Lett* 14:702–708
- Wansell SN, Geerts S, Coetzee JA (2022) Where are the seeds? Lack of floral morphs prevent seed production by the tristylous *Pontederia cordata* in South Africa. *Ecol Evol* 12:e9366
- Wilson JR (2024) A List of Taxa Currently and Historically Regulated under South Africa's National Environmental Management: Biodiversity Act, Alien & Invasive Species Regulations. doi: 10.5281/zenodo.8202704
- Wilson JR, Kumschick S (2024) The regulation of alien species in South Africa. *S Afr J Sci* 120, 10.17159/sajs.2024/17002
- Wilson RL, Hoch WA (2009) Identification of sterile, non-invasive cultivars of *Japanese spirea*. *HortScience* 44:2031–2034
- Wilson SB, Deng Z (2023) Ornamental Invasive Plants in Florida with Research-founded Alternatives. *HortTechnology* 33:349–356
- Wilson SB, Mecca LK (2003) Seed production and germination of eight cultivars and the wild type of *Ruellia tweediana*: a potentially invasive. *J Environ Hortic* 21:137–143
- Wirth FF, Davis KJ, Wilson SB (2004) Florida nursery sales and economic impacts of 14 potentially invasive landscape plant species. *J Environ Hortic* 22:12–16
- Zimmermann HG, Granata G (2002) Insect pests and diseases. Pages 235–254 in Nobel PS, ed. *Cacti: Biology and Uses*. Oakland: University of California Press

1 **Supplementary material:**

2 **Supplementary Table S1:** Unofficial consultative consensus list of “presumed sterile” cultivars and Hybrids (2014).

3 \*Number: this refers to the regulatory listing number as listed on the NEMBA 2014 Alien and Invasive Species regulatory list, where there were inconsistencies,  
4 numbers were changed accordingly.

5 \*The nomenclature for the species column was adjusted to accepted species names, and the genus and species names were removed to only show cultivar names  
6 in the final column.

7 \*NEMBA categories: **Category 1a** Species which must be combatted or eradicated; **Category 1b** Species which must be controlled.; **Category 2** listed species  
8 requires a permit issued by the Department of Forestry, Fisheries and the Environment (DFFE) to carry out a restricted activity; **Category 3** listed species are  
9 subject to certain exemptions in terms of section 70 (1)(a) of the NEMBA Act, which applies to the listing of alien invasive species. Any **Category 3** listed plant  
10 species that occurs in riparian areas, must be considered as **Category 1b** and the appropriate control measures instituted.

11 If there were no submission to the department “none” is flagged

Number	Species	Common name	NEMBA category	Sterile cultivars and hybrids
24	<i>Ageratum houstonianum</i>	Mexikaanse ageratum /Mexican ageratum	1b	F1 High Tide Blue   F1 High Tide White   F1 Blue Danube   F1 Blue Pearl   F1 Horizon Blue   F1 Blue Hawaii 5.0
56	<i>Berberis thunbergii</i>	Japanse berberis /Japanese barberry	3	Bagatelle   Golden King   Golden Ring   Green Ring   Harlequin   Kolbold   Nana   Ring of Fire   Rosy Glow   Ruby Gem
61	<i>Buddleja davidii</i>	Chinese sagewood, Summer Lilac	3	<i>Buddleja davidii</i> Hybrids: Flutterbye Grande Peach Cobbler   Flutterbye Grande Blueberry Cobbler   Flutterbye Grande Sweet marmalade   Flutterbye Grande Tangerine Dream   Flutterbye Grande Vanilla   Pink   Flutterbye Petite Snow White
69	<i>Callistemon/M elaleuca viminalis</i>	Treurperdestert /Weeping bottlebrush	1b in KwaZulu-Natal, Mpumalanga, Limpopo, Eastern Cape; 3 in the rest of South Africa	Burgundy Jack   Captain Cook   Endeavour   Hot Pink   Little John   Violaacea
78	<i>Catharanthus roseus</i>	Begraafplaasblom / Madagascar periwinkle	1b	Pacifica XP Mix   Pacifica XP Mystic Mix   Pacifica XP Orange   Pacifica XP Polka Dot   Pacifica XP Punch   Pacifica XP White   Pacifica XP Burgundy   Pacifica XP Burgundy   Halo   Pacifica XP Corral   Pacifica XP Cranberry   Pacifica XP Georgia Mix   Pacifica XP   Halo Mix   Pacifica XP Icy Pink   Pacifica XP Lilac   Pacifica XP Merlot Mix   Pacifica XP Raspberry   Pacifica XP Really Red   Pacifica XP Strawberry Shake Mix   Pacifica XP Cherry Red Halo   Mediterranean Deep Orchid   Mediterranean Deep Rose

Number	Species	Common name	NEMBA category	Sterile cultivars and hybrids
				Mediterranean Lilac   Mediterranean Polka Dot   Mediterranean XP Hot rose   Mediterranean XP Mix   Mediterranean XP Lipstick Mix   Mediterranean XP Peach   Mediterranean XP Red   Mediterranean XP Rose   Mediterranean XP Rose Halo   Mediterranean XP Dark red   Mediterranean XP Strawberry   Mediterranean XP White   Titan Apricot   Titan Blush   Titan Burgundy   Titan Halo Lavender Blue   Titan Lilac   Titan Mixture   Titan Polka Dot   Titan Punch   Titan Pure White   Titan Rose   Titan ROSE Halo   Titan Romance mix   Titan Bubble gum mix   Titan Clear Mix   Titan Cotton Candy mix   Titan Red white mix   Titan Romance Mix   Titan Summer Breeze Mix   Jams 'n Jellies   Blackberry Jams 'n Jellies   Blueberry Jams 'n Jellies American pie Mix   First Kiss Cherry   First Kiss Icy Pink   First Kiss Pink   First Kiss Polka Dot   First Kiss Raspberry
92	<i>Cestrum</i> spp.	Sestrum spp./Cestrum spp.	1b	none
99	<i>Coreopsis lanceolata</i>	Coreopsis /Tickseed	1a	none
101	<i>Cortaderia selloana</i>	Pampasgras, Silwergras / Pampas grass	1b	Silver Stripe   Gold Band
122	<i>Duranta erecta</i>	Vergeet-my-nie-boom/Forget-me-not-tree, Pigeon berry	3	Saphire Showers   Gold Mine   Sheena's Gold   Gold   Aussie   Iceberg
150	<i>Gleditsia triacanthos</i> <b>(Excluding cultivars under investigation)</b>	Amerikaanse drieroring/Honey locust	1b	Emerald Cascade   Inermis   Ruby Lace   Shademaster   Skyline   Sunburst
162	<i>Hedera canariensis</i>	Madeiraklimop/Canary ivy, Madeira ivy, Algerian ivy	3	Gloire de Marengo   Mosaik   Variegata
165	<i>Hedera helix</i>	Engelse hedera/English ivy	3	Buttercup   Chester   Crispa   Ferny   Golden hart   Jester Gold   Little Diamond   Marmorata   Golden child   Mini kolibri   Birgitte   Mini Wonder   Shamrock   Pittsburg   Tamara   White Ripple   Eva   Golden Cecilie   Mini White Wonder   Rochester
179	<i>Ipomoea indica</i>	Purperwinde/Morning glory	1b	none

Number	Species	Common name	NEMBA category	Sterile cultivars and hybrids
1178	<i>Ipomoea purpurea</i>	Purperwinde/Morning glory	1b	none
191	<i>Ligustrum lucidum</i>	Chinese blinkliguster/Chinese wax-leaved privet	1b in KwaZulu-Natal, Mpumalanga, North West, Limpopo, Eastern and Western Cape; 3 in Free State, Gauteng, Northern Cape	Tricolor
192	<i>Ligustrum ovalifolium</i>	Kaliforniese liguster/Californian privet	1b in KwaZulu Natal, Mpumalanga, North West, Limpopo, Eastern. Western Cape 3 in Free State, Gauteng, Northern Cape	Aurea
196	<i>Limonium sinuatum</i>	Statice, Sea Lavender	1b Northern Cape Western Cape	QIS Apricot   QIS Pale Blue   QIS Dark Blue   QIS Lavender   QIS Mix   QIS Pastel   QIS Rose   QIS White   QIS Yellow   Iceberg   Purple Attraction
212	<i>Metrosideros excelsa</i>	Nieu-Seelandse perdestert/New Zealand Christmas tree	1a in Overstrand District	Aurea   Maori Princess   Parnell   Spring Fire   Vibrance   Sunninghill
216	<i>Morus alba</i>	Witmoerbe, Gewone moerbe/White mulberry, Common mulberry	3	none
217	<i>Murraya paniculata</i>	Oranjejasmyn /Orange Jessamine	1b	Mini Mike
227	<i>Nephrolepis cordifolia</i>	Erect sword fern, Ladder sword fern	1b in KwaZulu Natal, Mpumalanga, Limpopo, Eastern; Western Cape; 3 in the rest of South Africa	Duffi   Kupukupu
228	<i>Nephrolepis exaltata</i> (L.) Schott (=	Swaardvaring /Sword fern, Boston sword fern	1b in KwaZulu-Natal, Mpumalanga, Limpopo, Eastern and	Boston Blue Bell   Boston   Corditas   Fluffy Ruffles   Green Delight   Green Moment   Marisa   Montana   Murano   Nevada   Teddy Junior?

Number	Species	Common name	NEMBA category	Sterile cultivars and hybrids
	<i>Polypodium exaltatum</i> L.)		Western Cape;3 in the rest of South Africa:	
229	<i>Nerium oleander</i> L.	Selonsroos /Oleander	1b	All double flowering cultivars   Petite Pink
239	<i>Opuntia ficus</i>	Boereturksvy, Grootdoringturksvy/ Mission prickly pear, Sweet prickly pear	1b	All spineless cactus pear cultivars and selections   cultivars grown for fruit production and fodder   Gymno Carpo   Roedtan   Morado   Tormentosa
245	<i>Opuntia robusta</i>	Robusta turksvy / Blue-leaf cactus	1a	none
265	<i>Pennisetum setaceum</i>	Pronkgras /Fountain grass	1b	<i>Pennisetum x advena</i> Rubrum   <i>Pennisetum x advena</i> Rubrum Fireworks   <i>Pennisetum x advena</i> Rubrum Skyrocket   All hybrids <i>P. setaceum x machrostachyum</i>   <i>Pennisetum</i> Vertigo
265	<i>Pyracantha angustifolia</i>	Geelbranddoring / Yellow firethorn	1b	none
299	<i>Pyracantha coccinea</i> <b>(Excluding cultivars under investigation)</b>	Rooibranddoring / Red firethorn	1b	none
300	<i>Pyracantha crenatoserrata</i>	Chinese firethorn, Broad leaf firethorn	1b	none
300	<i>Pyracantha crenulata</i>	Rooivuurddoring / Himalayan firethorn	1b	none
301	<i>Pyracantha koidzumii</i>	Formosa firethorn	1b	none
374	<i>Vinca major</i>	Gewone-opklim/Greater periwinkle	1b	Variegata   Blue Star
375	<i>Vinca minor</i>	Lesser periwinkle	1b	Alba   Multiplex   Illumination   Variegata

12 **Supplementary Table S2:** Unofficial consultative consensus list of “presumed sterile” cultivars and Hybrids (2022).

13 \*Number: this refers to the regulatory listing number as listed on the NEMBA 2014 Alien and Invasive Species regulatory list, where there were inconsistencies,  
14 numbers were changed accordingly.

15 \*The nomenclature for the species column was adjusted to accepted species names, and the genus and species names were removed to only show cultivar names  
16 in the final column.

17 \*NEMBA categories: **Category 1a** Species which must be combatted or eradicated; **Category 1b** Species which must be controlled.; **Category 2** listed species  
18 requires a permit issued by the Department of Forestry, Fisheries and the Environment (DFFE) to carry out a restricted activity; **Category 3** listed species are  
19 subject to certain exemptions in terms of section 70 (1)(a) of the NEMBA Act, which applies to the listing of alien invasive species. Any Category 3 listed plant  
20 species that occurs in riparian areas, must be considered as Category 1b and the appropriate control measures instituted.

21 If there were no submission to the department “none” is flagged

Number	Species	Common name	NEMBA category	Sterile cultivars and hybrids
24	<i>Ageratum houstonianum</i>	Mexikaanse ageratum /Mexican ageratum	1b	F1 High Tide Blue   F1 High Tide White   F1 Blue Danube   F1 Blue Pearl   F1 Horizon Blue   F1 Blue Hawaii 5.0
56	<i>Berberis thunbergii</i>	Japanse berberis /Japanese barberry	3	Bagatelle   Golden King   Golden Ring   Green Ring   Harlequin   Kolbold   Nana   Ring of Fire   Rosy Glow   Ruby Gem
69	<i>Callistemon/Melaleuca viminalis</i>	Treurperdestert /Weeping bottlebrush	1b in KwaZulu-Natal, Mpumalanga, Limpopo, Eastern Cape; 3 in the rest of South Africa	Burgundy Jack   Captain Cook   Endeavour   Hot Pink   Little John   Violaeca
78	<i>Catharanthus roseus</i>	Begraafplaasblom / Madagascar periwinkle	1b	Pacifica XP Mix   Pacifica XP Mystic Mix   Pacifica XP Orange   Pacifica XP Polka Dot   Pacifica XP Punch   Pacifica XP White   Pacifica XP Burgundy   Pacifica XP Burgundy   Halo   Pacifica XP Corral   Pacifica XP Cranberry   Pacifica XP Georgia Mix   Pacifica XP   Halo Mix   Pacifica XP Icy Pink   Pacifica XP Lilac   Pacifica XP Merlot Mix   Pacifica XP Raspberry   Pacifica XP Really Red   Pacifica XP Strawberry Shake Mix   Pacifica XP Cherry Red Halo   Mediterranean Deep Orchid   Mediterranean Deep Rose   Mediterranean Lilac   Mediterranean Polka Dot   Mediterranean XP Hot rose   Mediterranean XP Mix   Mediterranean XP Lipstick Mix   Mediterranean XP Peach   Mediterranean XP Red   Mediterranean XP Rose   Mediterranean XP Rose   Halo Mediterranean XP Dark red   Mediterranean XP Strawberry   Mediterranean XP White   Titan Apricot   Titan Blush   Titan Burgundy   Titan Halo Lavender Blue   Titan Lilac   Titan Mixture   Titan Polka Dot   Titan Punch   Titan Pure White   Titan Rose   Titan ROSE Halo   Titan Romance mix   Titan Bubble gum mix   Titan Clear Mix   Titan Cotton Candy mix   Titan Red white mix   Titan Romance Mix   Titan Summer Breeze Mix   Jams ‘n Jellies   Blackberry Jams ‘n Jellies   Blueberry Jams ‘n Jellies American pie Mix   First Kiss Cherry   First Kiss Icy Pink   First Kiss Pink   First Kiss Polka Dot   First Kiss Raspberry
122	<i>Duranta erecta</i>	Vergeet-my-nie-boom/Forget-me-not-tree, Pigeon berry	3	Saphire Showers   Gold Mine   Sheena’s Gold   Gold   Aussie   Iceberg

Number	Species	Common name	NEMBA category	Sterile cultivars and hybrids
162	<i>Hedera canariensis</i>	Madeiraklimop/Canary ivy, Madeira ivy, Algerian ivy	3	Gloire de Marengo   Mosaik   Variegata
165	<i>Hedera helix</i>	Engelse heder/English ivy	3	Buttercup   Chester   Crispa   Ferny   Golden hart   Jester Gold   Little Diamond   Marmorata   Golden child   Mini kolibri   Birgitte   Mini Wonder   Shamrock   Pittsburg   Tamara   White Ripple   Eva   Golden Cecilie   Mini White Wonder   Rochester Aurea
192	<i>Ligustrum ovalifolium</i>	Kaliforniese liguster/Californian privet	1b in KwaZuluNatal, Mpumalanga, North West, Limpopo, Eastern. Western Cape 3 in Free State, Gauteng, Northern Cape	
217	<i>Murraya paniculata</i>	Oranjejasmyn /Orange Jessamine	1b	Mini Mike
227	<i>Nephrolepis cordifolia</i>	Erect sword fern, Ladder sword fern	1b in KwaZuluNatal, Mpumalanga, Limpopo, Eastern; Western Cape; 3 in the rest of South Africa	Duffi   Kupukupu
228	<i>Nephrolepis exaltata</i> (L.) Schott (= <i>Polypodium exaltatum</i> L.)	Swaardvaring /Sword fern, Boston sword fern	1b in KwaZulu-Natal, Mpumalanga, Limpopo, Eastern and Western Cape; 3 in the rest of South Africa:	Boston Blue Bell   Boston   Corditas   Fluffy Ruffles   Green Delight   Green Moment   Marisa   Montana   Murano   Nevada   Teddy Junior'
239	<i>Opuntia ficus</i>	Boereturksvy, Grootdoringturksvy/ Mission prickly pear, Sweet prickly pear	1b	All spineless cactus pear cultivars and selections   cultivars grown for fruit production and fodder   Gymno Carpo   Roedtan   Morado   Tormentosa
245	<i>Opuntia robusta</i>	Robusta turksvy / Blue-leaf cactus	1a	none
265	<i>Pennisetum setaceum</i>	Pronkgras /Fountain grass	1b	<i>Pennisetum x advena</i> Rubrum   <i>Pennisetum x advena Rubrum</i> Fireworks   <i>Pennisetum x advena Rubrum</i> Skyrocket   All hybrids <i>P. setaceum x machrostachyum</i>   <i>Pennisetum</i> Vertigo
375	<i>Vinca minor</i>	Lesser periwinkle	1b	Alba   Multiplex   Illumination   Variegata

23 **Supplementary Appendix: Proposals by Government to the Green Industry (Information**  
24 **obtained from key stakeholder engagement meeting held on the 14th of February 2017)**

25 \* *The data on the original document was simplified by removing information such as regulatory name,*  
26 *common name, and NEMBA category (as this information was already included in Supplementary*  
27 *Table S1). In certain cases, text was adjusted from direct to indirect speech.*

28 1. The government should now require permits for the sale of the sterile cultivars and hybrids. This is  
29 because of the real difficulties that government are having in identifying the sterile hybrids and cultivars  
30 from the listed species. The government is not convinced that all nurseries and landscape architects are  
31 able to tell the differences either.

32 2. The following species should be removed from the lists of exempted species. *Coreopsis lanceolata*  
33 (tickseed); *Vinca major* (greater periwinkle). This is because tickseed is a hybrids or cultivars of a  
34 Category 1a species, and there is no proof of its sterility, and because the greater periwinkle is spreading  
35 more vegetatively than by seed, and it too must have a risk assessment before the exemption can be re-  
36 established:

37 3. The industry is given six months (from 14 February 2017) to provide the necessary proof of the  
38 sterility of the following cultivars and hybrids, failing which there shall be a withdrawal of the  
39 exemptions of sterile cultivars and hybrids:

40 *Ageratum houstonianum; Ipomoea indica; Ipomoea purpurea; Ligustrum lucidum; Ligustrum*  
41 *ovalifolium; Limonium sinuatum; Limonium sinuatum; Pyracantha angustifolia; Pyracantha coccinea;*  
42 *Pyracantha crenatoserrata; Pyracantha crenulata; Pyracantha koidzumii.*

43 4. The industry is given one year to provide the necessary proof of the sterility of the following cultivars  
44 and hybrids, failing which, the government shall withdraw the exemptions as sterile cultivars and  
45 hybrids:

46 *Callistemon viminalis; Canna indica; Catharanthus roseus; Cortaderia selloana; Gleditsia*  
47 *triacanthos; Metrosideros excelsa; Murraya paniculata; Nephrolepis cordifolia; Nerium oleander;*  
48 *Pennisetum setaceum; Vinca minor.*

49 5. The industry is given two years to provide the necessary proof of the sterility of the following  
50 cultivars and hybrids, failing which, the government shall withdraw the exemptions as sterile cultivars  
51 and hybrids:

52 *Acer negundo; Berberis thunbergia; Buddleja davidii; Cestrum* species (not specifically listed);  
53 *Duranta erecta; Hedera canariensis; Hedera helix; Morus alba.*

54 6. Note that *Pinus elliotti* is being changed to being only Category 2 in the upcoming amendments, and  
55 this is being addressed with the forestry industry.

56 7. It is also proposed that the government add the following nine species to the regulatory lists  
57 (following input from the Department of Agriculture, Forestry and Fisheries, and the Department of  
58 Water and Sanitation):

59 *Cynodon dactylon; Leptospermum scoparium; Lolium perenne; Lolium multiflorum; Morus nigra;*  
60 *Oenothera rosea; Oenothera stricta; Oenothera indecora.*